

Package leaflet: Information for the user

Fertin 1 mg tablets

Fertin 2 mg tablets

Fertin 3 mg tablets

Fertin 4 mg tablets

Glimepiride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Fertin is and what it is used for
2. What you need to know before you take Fertin
3. How to take Fertin
4. Possible side effects
5. How to store Fertin
6. Contents of the pack and other information

1. What Fertin is and what it is used for

Fertin is an orally active blood sugar lowering medicine. Fertin contains glimepiride, which belongs to a blood sugar lowering group of medicines called sulfonylurea. Fertin works by increasing the amount of insulin released from your pancreas. The insulin then lowers your blood sugar levels.

What Fertin is used for:

Fertin is used to treat a certain form of diabetes (type 2 diabetes mellitus) when diet, physical exercise and weight reduction alone have not been able to control your blood sugar levels.

2. What you need to know before you take Fertin

Do not take Fertin if you:

- are allergic to glimepiride or other sulfonylureas (medicines used to lower your blood sugar such as glibenclamide) or sulfonamides (medicines for bacterial infections such as sulfamethoxazole) or any of the other ingredients of this medicine (listed in section 6).
- have insulin dependent diabetes (type 1 diabetes mellitus).
- have diabetic ketoacidosis (a complication of diabetes when your acid level is raised in your body and you may have some of the following signs: fatigue, feeling sick (nausea), frequent urination and muscular stiffness).
- are in a diabetic coma.
- have severe kidney disease.
- have a severe liver disease.

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Fertin.

Warnings and precautions

Talk to your doctor or pharmacist before taking Fertin if you:

- are recovering from an injury, operation, infections with fever, or from other forms of stress, inform your doctor as temporary change of treatment may be necessary.
- have a severe liver or kidney disorder.

If you are not sure if any of these apply to you, talk to your doctor or pharmacist before taking Fertin.

Lowering of the haemoglobin level and breakdown of red blood cells (haemolytic anemia) can occur in patients missing the enzyme glucose-6-phosphate dehydrogenase.

The information available on the use of glimepiride in people under 18 years of age is limited. Therefore, its use in these patients is not recommended.

Important information about hypoglycaemia (low blood sugar)

When you take glimepiride, you may get hypoglycaemia (low blood sugar). Please see below for additional information about hypoglycaemia, its signs and treatment.

Following factors could increase the risk of you getting hypoglycaemia:

- Undernourishment, irregular meal time, missed or delayed meal or period of fasting
- Changes to your diet
- Taking more glimepiride than needed
- Having decreased kidney function
- Having severe liver disease
- If you suffer from particular hormone-induced disorders (disorders of the thyroid glands, of the pituitary gland or adrenal cortex)
- Drinking alcohol (especially when you skip a meal)
- Taking certain other medicines (see below "Other medicines and Fertin")
- If you increase your body exercise and you do not eat enough or eat food containing less carbohydrate than normal

Symptoms of hypoglycaemia include:

- Hunger pangs, headache, nausea, vomiting, sluggishness, sleepiness, disordered sleep, restlessness, aggression, impaired concentration, reduced alertness and reaction time, depression, confusion, speech and visual disorders, slurred speech, shakiness, partial paralysis, sensory disturbances, dizziness, helplessness.
- The following signs may also occur: sweating, clammy skin, anxiety, accelerated heart beat, high blood pressure, palpitations, sudden strong pain in the breast that may radiate into neighbouring areas (angina pectoris and cardiac arrhythmias).

If blood sugar levels continue to drop you may suffer from considerable confusion (delirium), develop convulsions, lose self control, breathing may be shallow and your heart beat slowed down, you may fall into unconsciousness. The clinical picture of a severe reduced blood sugar level may resemble that of a stroke.

Treating hypoglycaemia:

In most cases the signs of reduced blood sugar vanish very quickly when you consume some form of sugar, e.g. sugar cubes, sweet juice, sweetened tea.

You should therefore always take some form of sugar with you (e.g. sugar cubes). Remember that artificial sweeteners are not effective. Please contact your doctor or go to the hospital if taking sugar does not help or if the symptoms recur.

Laboratory tests

The level of sugar in your blood or urine should be checked regularly. Your doctor may also take blood tests to monitor your blood cell levels and liver function.

Children and adolescents:

Fertin is not recommended for use in children under 18 years of age.

Other medicines and Fertin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Your doctor may wish to change your dose of Fertin if you are taking other medicines, which may weaken or strengthen the effect of glimepiride on the level of sugar in your blood.

The following medicines can increase the blood sugar lowering effect of glimepiride. This can lead to a risk of hypoglycaemia (low blood sugar):

- Other medicines to treat diabetes mellitus (such as insulin or metformin)
- Medicines to treat pain and inflammation (phenylbutazone, azopropazone, oxyphenbutazone, aspirin-like medicines)
- Medicines to treat urinary infections (such as some long acting sulfonamides)
- Medicines to treat bacterial and fungal infections (tetracyclines, chloramphenicol, fluconazole, miconazole, quinolones, clarithromycin)
- Medicines to inhibit blood clotting (coumarin derivatives such as warfarin)
- Medicines supporting muscle build up (anabolics)
- Medicines used for male sex hormone replacement therapy
- Medicines to treat depression (fluoxetine, MAO-inhibitors)
- Medicines lowering high cholesterol level (fibrates)
- Medicines lowering high blood pressure (ACE inhibitors)
- Medicines called anti-arrhythmic agents used to control abnormal heart beat (disopyramide)
- Medicines to treat gout (allopurinol, probenecid, sulfinpyrazone)
- Medicines to treat cancer (cyclophosphamide, ifosfamide, trofosfamide)
- Medicines used to reduce weight (fenfluramine)
- Medicines to increase circulation when given in a high dose intravenous infusion (pentoxifylline)
- Medicines to treat nasal allergies such as hay fever (tritoqualine)
- Medicines called sympatholytics to treat high blood pressure, heart failure, or prostate symptoms

The following medicines may decrease the blood sugar lowering effect of glimepiride. This can lead to a risk of hyperglycaemia (high blood sugar level):

- Medicines containing female sex hormones (oestrogens, progestogens)
- Medicines supporting urine production (thiazide diuretics)
- Medicines used to stimulate the thyroid gland (such as levothyroxine)
- Medicines to treat allergies and inflammation (glucocorticoids)
- Medicines to treat severe mental disorders (chlorpromazine and other phenothiazine derivatives)
- Medicines used to raise heart beat, to treat asthma or nasal congestion, coughs and colds, used to reduce weight, or used in life-threatening emergencies (adrenaline and sympathomimetics)
- Medicines to treat high cholesterol level (nicotinic acid)
- Medicines to treat constipation when they are used long term (laxatives)
- Medicines to treat seizures (phenytoin)
- Medicines to treat nervousness and sleep problems (barbiturates)
- Medicines to treat increased pressure in the eye (azetazolamide)
- Medicines to treat high blood pressure or low blood sugar (diazoxide)
- Medicines to treat infections, tuberculosis (rifampicine)
- Medicines to treat severe low blood sugar levels (glucagon)

The following medicines can increase or decrease the blood sugar lowering effect of glimepiride:

- Medicines to treat stomach ulcers (called H2 antagonists)
- Medicines to treat high blood pressure or heart failure such as beta-blockers, clonidine, guanethidine and reserpine. These can also hide the signs of hypoglycaemia, so special care is needed when taking these medicines

Glimepiride may either increase or weaken the effects of the following medicines:

- Medicines inhibiting blood clotting (coumarin derivatives such as warfarin)

Colesevelam, a medicine used to reduce cholesterol, has an effect on the absorption of glimepiride. To avoid this effect, you should be advised to take Fertin at least 4 hours before colesevelam.

Fertin with food, drink and alcohol

Alcohol intake may increase or decrease the blood sugar lowering action of glimepiride in an unpredictable way.

Pregnancy, breastfeeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

Fertin should not be taken during pregnancy. Tell your doctor if you are, you think you might be or are planning to become pregnant.

Breast-feeding

Glimepiride may pass into breast milk. Fertin should not be taken during breast-feeding.

Driving and using machines

Your ability to concentrate or react may be impaired if your blood sugar is lowered (hypoglycaemia), or raised (hyperglycaemia) or if you develop visual problems as a result of such conditions. Bear in mind that you could endanger yourself or others (e.g. when driving a car or using machines). Please ask your doctor whether you can drive a car if you:

- have frequent episodes of hypoglycaemia,
- have fewer or no warning signals of hypoglycaemia

Fertin contains lactose monohydrate and sodium.

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium free'.

3. How to take Fertin

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Taking this medicine

- Take this medicine by mouth, just before or with the first main meal of the day (usually breakfast). If you do not have breakfast you should take the medicine on schedule as prescribed by your doctor. It is important not to leave out any meal when you are on Fertin.
- Swallow the tablets with at least half glass of water. Do not crush or chew the tablets.
- The tablets can be divided into equal doses.

How much to take

The dose of Fertin depends on your needs, condition and results of blood and urine sugar tests and is determined by your doctor. Do not take more tablets than your doctor has prescribed.

- The usual starting dose is one Fertin 1 mg tablet once a day.
- If necessary, your doctor may increase the dose after each 1 - 2 weeks of treatment.
- The maximum recommended dose is 6 mg Fertin per day.

- A combination therapy of glimepiride plus metformin or of glimepiride plus insulin may be initiated. In such a case your doctor will determine the proper doses of glimepiride, metformin or insulin individually for you.
- If your weight changes or if you change your lifestyle, or you are in a stress situation this may require changed Fertin doses, therefore inform your doctor.
- If you feel the effect of your medicine is too weak or too strong do not change the dose yourself, but ask your doctor.

If you take more Fertin than you should:

If you happen to have taken too much Fertin or an additional dose there is a danger of hypoglycaemia (signs of hypoglycaemia see section 2) and therefore you should instantly consume enough sugar (e.g. a small bar of sugar cubes, sweet juice, sweetened tea) and inform a doctor immediately. When treating hypoglycaemia due to accidental intake in children, the quantity of sugar given must be carefully controlled to avoid the possibility of producing dangerous hyperglycaemia. Persons in a state of unconsciousness must not be given food or drink.

Since the state of hypoglycaemia may last for some time it is very important that the patient is carefully monitored until there is no more danger. Admission into hospital may be necessary, also as a measure of precaution. Show the doctor the package or remaining tablets, so the doctor knows what has been taken.

Severe cases of hypoglycaemia accompanied by loss of consciousness and severe neurological failure are cases of medical emergency requiring immediate medical treatment and admission into hospital. It should be ensured that there is always a pre-informed person that can call a doctor in case of emergency.

If you forget to take Fertin:

If you forget to take a dose, do not take a double dose to make up for forgotten doses.

If you stop taking Fertin

If you interrupt or stop the treatment you should be aware that the desired blood sugar lowering effect is not achieved or that the disease will deteriorate again. Keep taking Fertin until your doctor tells you to stop.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects although not everybody gets them.

Tell your doctor immediately if you experience any of the following symptoms:

- Allergic reactions (including inflammation of blood vessels, often with skin rash) which may develop into serious reactions with difficulty in breathing, fall in blood pressure and sometimes progressing to shock.
- Abnormal liver function including yellowing of the skin and eyes (jaundice), problems with the bile flow (cholestasis), inflammation of the liver (hepatitis) or liver failure.
- Allergy (hypersensitivity) of the skin such as itching, rash, hives and increased sensitivity to sun. Some mild allergic reactions may develop into serious reactions.
- Severe hypoglycaemia including loss of consciousness, seizures or coma.

Some patients experienced the following side effects whilst taking glimepiride:

Rare side effects (may affect up to 1 in 1,000 people)

- Lower blood sugar than normal (hypoglycaemia) (see section 2).

- Decrease in the number of blood cells:
 - Blood platelets (which increases risk of bleeding or bruising).
 - White blood cells (which makes infections more likely).
 - Red blood cells (which can make the skin pale and cause weakness or breathlessness).
 These problems generally get better after you stop taking glimepiride.
- Weight gain
- Hair loss
- Taste disturbance.

Very rare side effects (may affect up to 1 in 10,000 people)

- Allergic reactions (including inflammation of blood vessels, often with skin rash) which may develop into serious reactions with difficulty in breathing, fall in blood pressure and sometimes progressing to shock. If you experience any of these symptoms, tell your doctor immediately.
- Abnormal liver function including yellowing of the skin and eyes (jaundice), impairment of the bile flow (cholestasis), inflammation of the liver (hepatitis) or liver failure. If you experience any of these symptoms, tell your doctor immediately.
- Feeling or being sick, diarrhoea, feeling full or bloated, and abdominal pain.
- Decrease in the amount of sodium level in your blood (shown by blood tests).

Other side effects (frequency not known – cannot be estimated from the available data) include:

- Allergy (hypersensitivity) of the skin may occur such as itching, rash, hives and increased sensitivity to sun. Some mild allergic reactions may develop into serious reactions with swallowing or breathing problems, swelling of your lips, throat or tongue. Therefore in the event of one of these side effects, tell your doctor immediately.
- Allergic reactions with sulfonylureas, sulfonamides, or related medicines may occur.
- Problems with your sight may occur when beginning treatment with glimepiride. This is due to changes in blood sugar levels and should soon improve.
- Increased liver enzymes.
- Severe unusual bleeding or bruising under the skin.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fertin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the package after EXP. The expiry date refers to the last day of that month.

This product does not require any special storage condition.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Fertin contains:

- The active ingredient is glimepiride. Each tablet contains 1mg, 2mg, 3mg or 4mg glimepiride.
- The other ingredients (excipients) are: Lactose monohydrate, Sodium starch glycolate, Povidone (K30), Magnesium stearate.

Fertin 1 mg tablet also contains the colouring agent E172 (red iron oxide), Fertin 2 mg contains E172 (yellow iron oxide) and E132 (indigo carmine), Fertin 3 mg contains E172 (yellow iron oxide), Fertin 4 mg contains E132 (indigo-carmine).

What Fertin looks like and contents of the pack

Fertin 1 mg is pink, round, flat, scored on one side tablet with diameter 6mm.

Fertin 2 mg is green, oval, flat, scored on one side, embossed “MC” on the other side tablet with dimensions 5x10mm.

Fertin 3 mg is pale yellow, oval, flat, scored on one side, embossed “MC” on the other side tablet with dimensions 5x10mm.

Fertin 4 mg is blue, oval, flat, scored on one side, embossed “MC” on the other side tablet with dimensions 5x10mm.

The score line helps dividing the tablets in two equal halves.

The tablets are packed in PVC/PVDC/Aluminum blisters. Packs of 30 tablets are available.

Marketing Authorisation Holder and Manufacturer

Medochemie Ltd, 1-10 Constantinoupoleos str, 3011 Limassol, Cyprus

This leaflet was last revised in June 2020.