

## Package leaflet: Information for the user

Bimato-Vision 0,1 mg/ml, oogdruppels, oplossing  
bimatoprost

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Bimato-Vision 0,1 mg/ml is and what it is used for
2. What you need to know before you use Bimato-Vision 0,1 mg/ml
3. How to use Bimato-Vision 0,1 mg/ml
4. Possible side effects
5. How to store Bimato-Vision 0,1 mg/ml
6. Contents of the pack and other information

### 1. What Bimato-Vision 0,1 mg/ml is and what it is used for

Bimato-Vision 0,1 mg/ml is an antiglaucoma preparation. It belongs to a group of medicines called prostamides.

Bimato-Vision 0,1 mg/ml is used to reduce high pressure in the eye. This medicine may be used on its own or with other drops called beta-blockers which also reduce pressure.

Your eye contains a clear, watery liquid that feeds the inside of the eye. Liquid is constantly being drained out of the eye and new liquid is made to replace this. If the liquid cannot drain out quickly enough, the pressure inside the eye builds up. This medicine works by increasing the amount of liquid that is drained. This reduces the pressure inside the eye. If the high pressure is not reduced, it could lead to a disease called glaucoma and eventually damage your sight.

### 2. What you need to know before you use Bimato-Vision 0,1 mg/ml

#### Do not use Bimato-Vision 0,1 mg/ml:

- if you are allergic to bimatoprost or any of the other ingredients of this medicine (listed in section 6).
- if you have had to stop using eye drops in the past because of a side effect of the preservative benzalkonium chloride.

#### Warnings and precautions

Talk to your doctor or pharmacist before using Bimato-Vision 0,1 mg/ml.

Talk to your doctor, if:

- You have any breathing problems.
- You have liver or kidney problems.

- You have had a cataract surgery in the past.
- You have dry eye.
- You have or have had any problems with your cornea (front transparent part of the eye).
- You wear contact lenses (see “X contains benzalkonium chloride”).
- You have or have had low blood pressure or low heart rate.
- You have had a viral infection or inflammation of the eye.

During treatment, Bimato-Vision 0,1 mg/ml may cause a loss of fat around the eye, which may cause your eyelid crease to deepen, your eye to appear sunken (enophthalmos), your upper eyelid to droop (ptosis), the skin around your eye to tighten (involution of dermatochalasis) and the lower white part of your eye to become more visible (inferior scleral show). The changes are typically mild, but if pronounced, they can affect your field of vision. The changes may disappear if you stop taking Bimato-Vision 0,1 mg/ml. Bimatoprost may also cause your eyelashes to darken and grow, and cause the skin around the eyelid to darken too. The colour of your iris may also go darker. These changes may be permanent. The change may be more noticeable if you are only treating one eye.

### **Children and adolescents**

Bimatoprost has not been tested in children under the age of 18 and therefore should not be used by patients under 18 years.

### **Other medicines and Bimato-Vision 0,1 mg/ml**

Tell your doctor or pharmacist if you are taking or have recently taken or might take any other medicines.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Bimato-Vision 0,1 mg/ml may get into breast milk so you should not breast-feed while you are taking Bimato-Vision 0,1 mg/ml.

### **Driving and using machines**

Your sight may become blurred for a short time just after using Bimato-Vision 0,1 mg/ml. You should not drive or use machines until your sight is clear again.

### **Bimato-Vision 0,1 mg/ml contains benzalkonium chloride**

This medicine contains 5 µg benzalkonium chloride in each drop (= 1 dose), which is equivalent to 0.20 mg/ml.

Benzalkonium chloride may be adsorbed by soft contact lenses and may change the colour of the contact lenses. You should remove contact lenses before using this medicine and put them back 15 minutes afterwards.

Benzalkonium chloride may also cause eye irritation, especially if you have dry eyes or disorders of the cornea (the clear layer at the front of the eye). If you feel abnormal eye sensation, stinging or pain in the eye after using this medicine, talk to your doctor.

### **Bimato-Vision 0,1 mg/ml contains phosphates**

This medicine contains 23.7 µg phosphates in each drop (= 1 dose), which is equivalent to 0.95 mg/ml. If you suffer from severe damage to the clear layer at the front of the eye (the cornea), phosphates may cause in very rare cases cloudy patches on the cornea due to calcium build-up during treatment.

### 3. How to use Bimato-Vision 0,1 mg/ml

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.


Bimato-Vision 0,1 mg/ml should only be applied to the eye. The recommended dose is one drop of Bimato-Vision 0,1 mg/ml in the evening, once daily in each eye that needs treatment.

If you use Bimato-Vision 0,1 mg/ml with another eye medicine, wait at least five minutes between using Bimato-Vision 0,1 mg/ml and the other eye medicine.

Do not use more than once a day as the effectiveness of treatment may be reduced.

#### Instructions for use:

You must not use the bottle if the tamper-proof seal on the bottle neck is broken before you first use it.



1. Wash your hands. Tilt your head back and look at the ceiling.
2. Gently pull down the lower eyelid until there is a small pocket.
3. Turn the bottle upside down and squeeze it to release one drop into each eye that needs treatment.
4. Let go of the lower lid, and close your eye for 30 seconds.  
Wipe off any excess that runs down the cheek.  
If a drop misses your eye, try again.

To help prevent infections and avoid eye injury, do not let the tip of the bottle touch your eye or anything else. Put the cap back on and close the bottle straight after you have used it.

#### If you use more Bimato-Vision 0,1 mg/ml than you should

If you use more Bimato-Vision 0,1 mg/ml than you should, it is unlikely to cause you any serious harm. Put your next dose in at the usual time. If you are worried, talk to your doctor or pharmacist.

#### If you forget to use Bimato-Vision 0,1 mg/ml

If you forget to use Bimato-Vision 0,1 mg/ml, use a single drop as soon as you remember, and then go back to your regular routine. Do not take a double dose to make up for a forgotten dose.

#### If you stop using Bimato-Vision 0,1 mg/ml

Bimato-Vision 0,1 mg/ml should be used every day to work properly. If you stop using Bimato-Vision 0,1 mg/ml the pressure inside your eye may go up, therefore talk to your doctor before stopping this treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Very common: may affect more than 1 in 10 people**

Affecting the eye

- Slight redness (up to 29 % of people)

Affecting the eye region:

- Loss of fat in the eye region which can lead to deepening of your eyelid crease, sunken eye (enophthalmos), drooping eyelid (ptosis), tightening of the skin around your eye (involution of dermatochalasis), and the lower white part of your eye to become more visible (inferior scleral show).

**Common: may affect up to 1 in 10 people**

Affecting the eye

- Small breaks in the surface of the eye, with or without inflammation
- Irritation
- Itchy eyes
- Longer eyelashes
- Irritation, when drop is put in the eye
- Eye pain

Affecting the skin

- Red and itchy eyelids
- Darker skin colour around the eye
- Hair growth around the eye

**Uncommon: may affect up to 1 in 100 people**

Affecting the eye

- Darker Iris colour
- Tired eye
- Swelling of the surface of the eye
- Blurred vision
- Loss of eye lashes

Affecting the skin

- Dry skin
- Crusting on the edge of the eyelid
- Swelling of the eyelid
- Itching

Affecting the body

- Headache
- Feeling of sickness

**Not known: frequency cannot be estimated from the available data**

Affecting the eye

- Macular oedema (swelling of the retina at the back of the eye which may lead to worsening vision)
- Darker eyelid colour
- Eye dryness

- Ocular discomfort
- Sensitivity to light
- Sticky eyes
- A feeling that something is in your eye
- Swelling of the eye
- Increasing tears

#### Affecting the body

- Symptoms of allergic reaction (swelling, redness of the eye and rash of the skin)
- Asthma
- Worsening of asthma
- Worsening of the lung disease called chronic obstructive pulmonary disease (COPD)
- Shortness of breath
- Dizziness
- Increased blood pressure
- Skin discoloration (periocular)

In addition to the side effects for Bimato-Vision 0,1 mg/ml, the following side effects have been seen with another medicine containing a higher strength of bimatoprost (0.3 mg/ml):

- Ocular burning
- An allergic reaction in the eye
- Inflamed eyelids
- Difficulty in seeing clearly
- Worsening of vision
- Swelling of the see-through layer that covers the eye
- A feeling that something is in your eye
- Tears
- Sticky eyes
- Darker eyelashes
- Retinal bleeding
- Inflammation within the eye;
- Cystoid macular oedema (swelling of the retina within the eye leading to worsening vision)
- Eyelid twitching
- Eyelid shrinking, moving away from surface of the eye
- Skin redness around the eye
- Weakness
- An increase in blood-test results that show how your liver is working.

#### **Other side effects reported with eye drops containing phosphates**

If you suffer from severe damage to the clear layer at the front of the eye (the cornea), phosphates may cause in very rare cases cloudy patches on the cornea due to calcium build-up during treatment.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the national reporting system

Nederland  
Nederlands Bijwerkingen Centrum Lareb  
Website: [www.lareb.nl](http://www.lareb.nl)

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Bimato-Vision 0,1 mg/ml**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle label and the carton after “EXP:”. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

You must throw away the bottle at the latest four weeks after you first opened it, even if there are still some drops left. This will prevent infections. To help you remember, write down the date you opened it in the space on the box.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Bimato-Vision 0,1 mg/ml contains**

- The active substance is bimatoprost. One ml of solution contains 0.1 mg bimatoprost. One drop contains approximately 2.5 micrograms bimatoprost.
- The other ingredients are benzalkonium chloride (preservative), sodium chloride, disodium phosphate heptahydrate, citric acid monohydrate, sodium hydroxide or hydrochloric acid (for pH-adjustment) and purified water.

### **What Bimato-Vision 0,1 mg/ml looks like and contents of the pack**

Bimato-Vision 0,1 mg/ml is a clear, colourless solution in a pack containing either 1 plastic bottle or 3 plastic bottles each with a screw cap. Each bottle is approximately half full and contains either 2.5 milliliters or 3 milliliters of solution. This is enough for 4 weeks' usage.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

OmniVision GmbH  
Lindberghstr. 9  
82178 Puchheim  
Duitsland

**Manufacturer**

Rompharm Company S.R.L.

Eroilor Street, no. 1A

Otopeni 075100, Ilfov

Roemenie

Farmigea S.p.A.

Via Giovan Battista Oliva 6-8

Pisa, 56121

Italië

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Nederland	Bimato-Vision 0,1 mg/ml, oogdruppels, oplossing
Duitsland	Bimato-Vision 0,1 mg/ml Augentropfen, Lösung
Oostenrijk	Bimato-Vision 100 Mikrogramm/ml Augentropfen, Lösung

**Deze bijsluiter is voor het laatst goedgekeurd in december 2024**