

## **Package leaflet: Information for the user**

### **Ulipristalacetaat Exeltis 30 mg film-coated tablet**

Ulipristal acetate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

Always take this medicine exactly as described in this leaflet or as your pharmacist, doctor or other healthcare professional has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your pharmacist, doctor, or other healthcare professional. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Ulipristalacetaat Exeltis is and what it is used for
2. What you need to know before you use Ulipristalacetaat Exeltis
3. How to use Ulipristalacetaat Exeltis
4. Possible side effects
5. How to store Ulipristalacetaat Exeltis
6. Contents of the pack and other information
  - Useful information about contraception

#### **1. What Ulipristalacetaat Exeltis is and what it is used for**

##### Ulipristalacetaat Exeltis is an emergency contraceptive

Ulipristalacetaat Exeltis is a contraceptive intended to prevent pregnancy after unprotected sex or if your contraceptive method has failed. For example:

- if you had sex without protection;
- if your or your partner's condom tore, slipped or came off, or if you forgot to use one;
- if you did not take your contraceptive pill as recommended.

You should take the tablet as soon as possible after sex, and within a maximum of 5 days (120 hours).

This is because the sperm can survive up to 5 days in your body after intercourse.

This medicine is suitable for any woman of childbearing age, including adolescents.

You can take this tablet at any time in the menstrual cycle.

##### Ulipristalacetaat Exeltis does not work if you are already pregnant.

If your menstrual period is late, there is a possibility that you may be pregnant. When your period is late or when you have symptoms of pregnancy (heavy breasts, morning sickness) you should consult a doctor or other healthcare professional before taking the tablet.

If you have unprotected sex after taking the tablet, it will not stop you from becoming pregnant. Unprotected sex at any time during your cycle can lead to pregnancy.

### Ulipristalacetaat Exeltis is not to be used for regular contraception

If you do not have a regular method of contraception, talk to your doctor or healthcare professional to choose one that is suitable for you.

### How Ulipristalacetaat Exeltis works

Ulipristalacetaat Exeltis contains the substance *ulipristal acetate* which acts by modifying the activity of the natural hormone progesterone which is necessary for ovulation to occur. As a result, this medicine works by postponing ovulation. Emergency contraception is not effective in every case. Of 100 women who take this medicine approximately 2 will become pregnant.

This medicine is a contraceptive used to prevent a pregnancy from starting. If you are already pregnant, it will not interrupt an existing pregnancy.

### Emergency contraception does not protect against sexually transmitted infections

Only condoms can protect you from sexually transmitted infections. This medicine will not protect you against HIV infection or any other sexually transmitted diseases (e.g. chlamydia, genital herpes, genital warts, gonorrhoea, hepatitis B and syphilis). Ask a healthcare professional for advice if you are worried about this.

There is more information about contraception at the end of this leaflet.

## **2. What you need to know before you take Ulipristalacetaat Exeltis**

### **Do not take Ulipristalacetaat Exeltis**

- If you are allergic to ulipristal acetate or any of the other ingredients of this medicine (listed in section 6).

### **Warning and precautions**

Talk to your pharmacist, doctor or other healthcare professional before taking Ulipristalacetaat Exeltis

- if your period is late or you have symptoms of pregnancy (heavy breasts, morning sickness), as you may already be pregnant (see section “Pregnancy, breast-feeding and fertility”);
- if you suffer from severe asthma;
- if you suffer from severe liver disease.

In all women, emergency contraception should be taken as soon as possible after unprotected intercourse. There is some evidence that this medicine may be less effective with increasing body weight or body mass index (BMI), but these data were limited and inconclusive. Therefore, Ulipristalacetaat Exeltis is still recommended for all women regardless of their weight or BMI.

You are advised to speak to a healthcare professional if you are concerned about any problems related to taking emergency contraception.

If you become pregnant despite taking this medicine, it is important that you see your doctor. See section “Pregnancy, breast-feeding and fertility” for more information.

## **Other contraceptives and Ulipristalacetaat Exeltis**

This medicine may make regular hormonal contraceptives, like pills and patches, temporarily less effective.

If you are currently taking hormonal contraception, continue to use it as usual after taking the tablet, but be sure to use condoms every time you have sex until your next period.

Do not use Ulipristalacetaat Exeltis together with another emergency contraceptive pill that contains levonorgestrel.

By taking them both together, you might make this medicine less effective.

## **Other medicines and Ulipristalacetaat Exeltis**

Tell your pharmacist, doctor or other healthcare professional if you are taking or have recently taken any other medicines, including medicines obtained without a prescription or herbal medicines.

Some medicines may prevent Ulipristalacetaat Exeltis from working effectively. If you have used any of the medicines listed below during the last 4 weeks, Ulipristalacetaat Exeltis may be less suitable for you. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD):

- medicines used to treat epilepsy (for example, primidone, phenobarbital, phenytoin, fosphenytoine, carbamazepine, oxcarbazepine and barbiturates)
- medicines used to treat tuberculosis (for example, rifampicin, rifabutin)
- a treatment for HIV (ritonavir, efavirenz, nevirapine)
- a medicine used to treat fungal infections (griseofulvin)
- herbal remedies containing St John's wort (*Hypericum perforatum*).

Speak to your doctor or pharmacist before using Ulipristalacetaat Exeltis when you use (or have recently used) any of the medicines stated above.

## **Pregnancy, breast-feeding and fertility**

### Pregnancy

Before taking this medicine, if your period is late, tell your pharmacist, doctor or other healthcare professional, or do a pregnancy test in order to make sure you are not already pregnant (see section "Warning and precautions").

This medicine is a contraceptive used to prevent a pregnancy from starting. If you are already pregnant it will not interrupt an existing pregnancy.

"If you become pregnant despite taking this medicine, there is no evidence that it will affect your pregnancy. However, it is important that you see your doctor. As for any pregnancy, your doctor may want to check that the pregnancy is not outside the womb. This is especially important if you have severe abdominal (stomach) pain or bleeding or if you have previously had a pregnancy outside the womb, tubal surgery or long term (chronic) genital infection."

If you become pregnant despite taking Ulipristalacetaat Exeltis, you are encouraged to ask your doctor to register your pregnancy in an official registry. You can also report this information on your own at [www.ulipristal-pregnancy-registry.com](http://www.ulipristal-pregnancy-registry.com). Your information will remain anonymous – nobody will know it is information about you. Sharing your information may help women in the future understand the safety or risks of Ulipristalacetaat Exeltis during a pregnancy.

### Breast-feeding

If you take this medicine while you are breastfeeding a baby, do not breast-feed for one week after taking Ulipristalacetaat Exeltis. During this time, it is recommended to use a breast pump in order to maintain milk production, but throw away your breast milk. The effect of breastfeeding your baby in the week after taking this medicine is not known.

### Fertility

This medicine will not affect your future fertility. If you have unprotected sex after taking this medicine, it will not stop you from becoming pregnant. Therefore it is important you use condoms until your next period.

If you wish to start or continue with a regular method of contraception after using this medicine, you can do so but you should also use condoms until your next period.

### **Driving and using machines**

After taking this medicine, some women experience dizziness, drowsiness, blurred vision and/or loss of concentration (see section 4 “Possible side effects”). If you experience these symptoms, do not drive or use machines.

### **Ulipristalacetaat Exeltis contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

### **Ulipristalacetaat Exeltis contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

## **3. How to take Ulipristalacetaat Exeltis**

Always take this medicine exactly as described in this leaflet or as your pharmacist, doctor or other healthcare professional has told you. Check with your pharmacist or doctor if you are not sure.

### How to take the Ulipristalacetaat Exeltis film-coated tablet

- Take one tablet by mouth as soon as possible and no later than 5 days (120 hours) after unprotected sex or contraceptive failure. Take the tablet without delay.
- You can take this medicine at any time in your cycle.
- You can take this medicine at any time of the day either before, during or after a meal.
- If you are using one of the medicines that may prevent Ulipristalacetaat Exeltis from working properly (see section above “Other medicines and Ulipristalacetaat Exeltis”) or if you have used one of these medicines in the past 4 weeks, Ulipristalacetaat Exeltis may work less effectively for you. Speak to your doctor or pharmacist before using Ulipristalacetaat Exeltis. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD).

### If you vomit after taking Ulipristalacetaat Exeltis

If you vomit (be sick, throw up) within 3 hours of taking the tablet, take another tablet as soon as possible.

#### If you have sex again after taking Ulipristalacetaat Exeltis

If you have unprotected sex after taking the tablet, it will not stop you from becoming pregnant. After you take the tablet and until your next period comes, you should use condoms every time you have sex.

#### If your next period is late after taking Ulipristalacetaat Exeltis

After taking the tablet, it is normal for your next period to be a few days late. However, if your period is more than 7 days late; if it is unusually light or unusually heavy; or if you experience symptoms such as abdominal (stomach) pain, breast tenderness, vomiting or nausea, you may be pregnant. You should do a pregnancy test right away. If you are pregnant, it is important that you see your doctor. (See section “Pregnancy, breast-feeding and fertility”)

#### If you take more Ulipristalacetaat Exeltis than you should

There have been no reports of harmful effects from taking a higher dose than recommended of this medicine. However, do ask your pharmacist, doctor or other healthcare professional for advice. If you have any further questions on the use of this medicine, ask your pharmacist, doctor or other healthcare professional.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some symptoms such as breast tenderness and abdominal (stomach) pain, throwing up (vomiting), feeling sick (nausea) are also possible signs of pregnancy. If you miss your period and experience such symptoms after taking Ulipristalacetaat Exeltis, you should do a pregnancy test (See section “Pregnancy, breastfeeding and fertility”).

#### Common side effects (may affect up to 1 in 10 people)

- nausea, abdominal (stomach) pain or discomfort, vomiting
- painful periods, pelvic pain, breast tenderness
- headache, dizziness, mood swings
- muscle pain, back pain, tiredness

#### Uncommon side effects (may affect up to 1 in 100 people)

- diarrhoea, heartburn, wind, dry mouth
- unusual or irregular vaginal bleeding, heavy/prolonged periods, premenstrual syndrome, vaginal irritation or discharge, lesser or greater sex drive
- hot flushes
- appetite changes, emotional disorders, anxiety, agitation, trouble sleeping, sleepiness, migraine, visual disturbances
- influenza
- acne, skin lesions, itching
- fever, chills, malaise

Rare side effects (may affect up to 1 in 1,000 people)

- allergic reactions such as rash, hives or swelling of the face
- genital pain or itching, pain during sex, rupture of an ovarian cyst, unusually light period
- loss of concentration, vertigo, shaking, disorientation, fainting
- unusual sensation in eye, red eye, sensitivity to light
- dry throat, disturbance in taste
- feeling thirsty

#### Reporting of side effects

If you get any side effects, talk to your pharmacist, doctor or other healthcare professional. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can provide more information on the safety of this medicine.

### **5. How to store Ulipristalacetaat Exeltis**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is on the carton and on the blister after marked EXP. The expiry date refers to the last day of that month.

Store in the original package in order to protect from light.

This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via waste water. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Ulipristalacetaat Exeltis contains**

- The active substance is ulipristal acetate. One film-coated tablet contains 30 milligrams of ulipristal acetate.
- The other ingredients are lactose monohydrate, pregelatinised starch (Maize), sodium starch glycolate, magnesium stearate, hypromellose (E464), hydroxypropylcellulose (E463), stearic acid (E570), talc (E553b), titanium dioxide (E171).

#### **What Ulipristalacetaat Exeltis looks like and contents of the pack**

Ulipristalacetaat Exeltis is a white, round, biconvex film-coated tablet of 9.0 - 9.2 mm diameter embossed with "U30" on one side.

Ulipristalacetaat Exeltis is available in carton containing one transparent PVC/PVDC/aluminium blister of 1 tablet.

**Marketing Authorisation Holder**

Exeltis Healthcare S.L  
Av.Miralcampo 7-Poligono  
Ind.Miralcampo| 19200,  
Azuqueca de Henares  
Guadalajara,  
Spain

**Manufacturer**

Haupt Pharma Münster GmbH  
Schleebrüggenkamp 15  
48159 Münster  
Germany

**This medicinal product is authorised in the Member States of the EEA under the following names:**

The Netherlands	Ulipristalacetaat Exeltis 30 mg, Filmomhulde tablet
France	Ulipristal Acétate Exeltis 30 mg, comprimé pelliculé

**Deze bijsluiter is voor het laatst goedgekeurd in september 2023**

## USEFUL INFORMATION ABOUT CONTRACEPTION

### MORE ABOUT EMERGENCY CONTRACEPTION

The sooner you take emergency contraception, the better the chance of avoiding pregnancy. Emergency contraception will not affect your fertility.

Emergency contraception can delay ovulation within a given menstrual cycle, but it will not stop you from becoming pregnant if you have unprotected sex again. After you take emergency contraception and until your next period comes, you should use a condom every time you have sex.

### MORE ABOUT REGULAR CONTRACEPTION

If you have taken emergency contraception and you do not use a regular contraceptive method (or do not have a contraceptive method that suits you), talk to your doctor or family planning clinic for advice.

There are many different types of contraception available, and you should be able to find the right method for you.

Examples of regular contraception methods:

---

#### Daily methods

Contraceptive pill

---

#### Weekly or monthly methods

Contraceptive patch

Vaginal Ring

---

#### Long lasting methods

Contraceptive implant

IUD (intrauterine device)

---