

Package leaflet: Information for the user

Capecitabine Sandoz® 150 mg, filmomhulde tabletten Capecitabine Sandoz® 500 mg, filmomhulde tabletten

capecitabine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Nationally completed name] is and what it is used for
2. What you need to know before you take [Nationally completed name]
3. How to take [Nationally completed name]
4. Possible side effects
5. How to store [Nationally completed name]
6. Contents of the pack and other information

1. What [Nationally completed name] is and what it is used for

[Nationally completed name] belongs to the group of medicines called "cytostatic medicines", which stop the growth of cancer cells. [Nationally completed name] contains capecitabine, which itself is not a cytostatic medicine. Only after being absorbed by the body is it changed into an active anti-cancer medicine (more in tumour tissue than in normal tissue).

[Nationally completed name] is used in the treatment of colon, rectal, gastric, or breast cancers. Furthermore, [Nationally completed name] is used to prevent new occurrence of colon cancer after complete removal of the tumour by surgery.

[Nationally completed name] may be used either alone or in combination with other medicines.

2. What you need to know before you take [Nationally completed name]

Do not take [Nationally completed name]:

- if you are allergic to capecitabine or any of the other ingredients of this medicine (listed in section 6). You must inform your doctor if you know that you have an allergy or over-reaction to this medicine,
- if you previously have had severe reactions to fluoropyrimidine therapy (a group of anticancer medicines such as fluorouracil),
- if you are pregnant or breast-feeding,
- if you have severely low levels of white cells or platelets in the blood (leucopenia, neutropenia or thrombocytopenia),
- if you have severe liver or kidney problems,
- if you know that you do not have any activity of the enzyme dihydropyrimidine dehydrogenase (DPD) (complete DPD deficiency)

- if you are being treated now or have been treated in the last 4 weeks with brivudine as part of herpes zoster (chickenpox or shingles) therapy.

Warnings and precautions

Talk to your doctor or pharmacist before taking [Nationally completed name] if you

- know that you have a partial deficiency in the activity of the enzyme dihydropyrimidine dehydrogenase (DPD)
- have a family member who has partial or complete deficiency of the enzyme dihydropyrimidine dehydrogenase (DPD)
- have liver or kidney diseases
- have or had heart problems (for example an irregular heartbeat or pains to the chest jaw and back brought on by physical effort and due to problems with the blood flow to the heart)
- have brain diseases (for example, cancer that has spread to the brain, or nerve damage (neuropathy))
- have calcium imbalances (seen in blood tests)
- have diabetes
- cannot keep food or water in your body because of severe nausea and vomiting
- have diarrhoea
- are or become dehydrated
- have imbalances of ions in your blood (electrolyte imbalances, seen in tests)
- have a history of eye problems as you may need extra monitoring of your eyes
- have a severe skin reaction.

DPD deficiency: DPD deficiency is a genetic condition that is not usually associated with health problems unless you receive certain medicines. If you have DPD deficiency and [Nationally completed name], you are at an increased risk of severe side effects (listed under section 4 Possible side effects). It is recommended to test you for DPD deficiency before start of treatment. If you have no activity of the enzyme you should not take [Nationally completed name]. If you have a reduced enzyme activity (partial deficiency) your doctor might prescribe a reduced dose. If you have negative test results for DPD deficiency, severe and life-threatening side effects may still occur.

Children and adolescents

[Nationally completed name] is not indicated in children and adolescent. Do not give [Nationally completed name] to children and adolescent.

Other medicines and [Nationally completed name]

Before starting treatment, tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is extremely important, as taking more than one medicine at the same time can strengthen or weaken the effect of the medicines.

**You must not take brivudine (an anti-viral medicine for the treatment of shingles or chickenpox) at the same time as capecitabine treatment (including during any rest periods when you are not taking any capecitabine tablets).
If you have taken brivudine you must wait for at least 4 weeks after stopping brivudine before starting to take capecitabine. See also section “Do not take [Nationally completed name]”.**

Also, you need to be particularly careful if you are taking any of the following:

- gout medicines (allopurinol),
- blood-thinning medicines (coumarin, warfarin),
- medicines for seizures or tremors (phenytoin)
- interferon alpha
- radiotherapy and certain medicines used to treat cancer (folinic acid, oxaliplatin, bevacizumab, cisplatin, irinotecan),
- medicines used to treat folic acid deficiency.

[Nationally completed name] with food and drink

You should take [Nationally completed name] no later than 30 minutes after meals.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. You must not take [Nationally completed name] if you are pregnant or think you might be. You must not breast-feed if you are taking [Nationally completed name] and for 2 weeks after the last dose. If you are a woman who could become pregnant you should use effective contraception during treatment with [Nationally completed name] and for 6 months after the last dose. If you are a male patient and your female partner could become pregnant, you should use effective contraception during treatment with [Nationally completed name] and for 3 months after the last dose.

Driving and using machines

[Nationally completed name] may make you feel dizzy, nauseous or tired. It is therefore possible that [Nationally completed name] could affect your ability to drive a car or operate machinery.

[Nationally completed name] contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

[Nationally completed name] contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take [Nationally completed name]

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

[Nationally completed name] should only be prescribed by a doctor experienced in the use of anticancer medicines.

Your doctor will prescribe a dose and treatment regimen that is right for *you*. The dose of [Nationally completed name] is based on your body surface area. This is calculated from your height and weight. The usual dose for adults is 1250 mg/m² of body surface area taken two times daily (morning and evening). Two examples are provided here: A person whose body weight is 64 kg and height is 1.64 m has a body surface area of 1.7 m² and should take 4 tablets of 500 mg and 1 tablet of 150 mg two times daily. A person whose body weight is 80 kg and height is 1.80 m has a body surface area of 2.00 m² and should take 5 tablets of 500 mg two times daily.

Your doctor will tell you what dose you need to take, when to take it and for how long you need to take it.

Your doctor may want you to take a combination of 150 mg and 500 mg tablets for each dose.

- Take the tablets **morning and evening as** prescribed by your doctor.
- Take the tablets within **30 minutes after the end of a meal** (breakfast and dinner) and swallow whole with water. Do not crush or cut tablets. If you cannot swallow [Nationally completed name] tablets whole, tell your healthcare provider.

- It is important that you take all your medicine as prescribed by your doctor.

[Nationally completed name] tablets are usually taken for 14 days followed by a 7 day rest period (when no tablets are taken). This 21 day period is one treatment cycle.

In combination with other medicines the usual dose for adults may be less than 1250 mg/m^2 of body surface area, and you may need to take the tablets over a different time period (e.g. every day, with no rest period).

If you take more [Nationally completed name] than you should

If you take more [Nationally completed name] than you should, contact your doctor as soon as possible before taking the next dose.

You might get the following side effects if you take a lot more capecitabine than you should: feeling or being sick, diarrhoea, inflammation or ulceration of the gut or mouth, pain or bleeding from the intestine or stomach, or bone marrow depression (reduction in certain kinds of blood cells). Tell your doctor immediately if you experience any of these symptoms.

If you forget to take [Nationally completed name]

Do not take the missed dose at all. Do not take a double dose to make up for a forgotten dose. Instead, continue your regular dosing schedule and check with your doctor.

If you stop taking [Nationally completed name]

There are no side-effects caused by stopping treatment with [Nationally completed name]. In case you are using coumarin anticoagulants (containing e.g. phenprocoumon), stopping [Nationally completed name] might require that your doctor adjusts your anticoagulant dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking [Nationally completed name] immediately and contact your doctor if any of these symptoms occur:

- **Diarrhoea:** if you have an increase of 4 or more bowel movements compared to your normal bowel movements each day or any diarrhoea at night.
- **Vomiting:** if you vomit more than once in a 24-hour time period.
- **Nausea:** if you lose your appetite, and the amount of food you eat each day is much less than usual.
- **Stomatitis:** if you have pain, redness, swelling or sores in your mouth and/or throat.
- **Hand-and-foot skin-reaction:** if you have pain, swelling, redness or tingling of hands and/or feet.
- **Fever:** if you have a temperature of 38°C or greater
- **Infection:** if you experience signs of infection caused by bacteria or virus, or other organisms.
- **Chest pain:** if you experience pain localised to the centre of the chest, especially if it occurs during exercise.
- **Steven-Johnson syndrome:** if you experience painful red or purplish rash that spreads and blisters and/or other lesions begin to appear in the mucous membrane (e.g. mouth and lips), in particular if you had before light sensitivity, infections of the respiratory system (e.g. bronchitis) and/or fever.
- **DPD Deficiency:** if you have a known DPD deficiency, you are at an increased risk of acute early-onset of toxicity and severe, life-threatening, or fatal adverse reactions caused by capecitabine (e.g. stomatitis, mucosal inflammation, diarrhoea, neutropenia, and neurotoxicity)
- **Angioedema:** Seek medical attention straight away if you notice any of the following symptoms -

you may need urgent medical treatment: swelling mainly of the face, lips, tongue or throat which makes it difficult to swallow or breathe, itching and rashes. This could be a sign of angioedema.

If caught early, these side effects usually improve within 2 to 3 days after treatment discontinuation. If these side effects continue, however, contact your doctor immediately. Your doctor may instruct you to restart treatment at a lower dose.

If severe stomatitis (sores in your mouth and/or throat), mucosal inflammation, diarrhoea, neutropenia (increased risk for infections), or neurotoxicity occurs during the first cycle of treatment a DPD deficiency may be involved (please see Section 2: Warning and precautions).

Hand and foot skin-reaction can lead to loss of fingerprint, which could impact your identification by fingerprint scan.

In addition to the above, when [Nationally completed name] is used alone, very common side effects which may affect more than 1 in 10 people are:

- abdominal pain
- rash, dry or itchy skin
- tiredness
- loss of appetite (anorexia)

These side effects can become severe; therefore, it is important that you **always contact your doctor immediately** when you start to experience a side effect. Your doctor may instruct you to decrease the dose and/or temporarily discontinue treatment with [Nationally completed name]. This will help reduce the likelihood that the side effect continues or becomes severe.

Other side effects are:

Common side effects (may affect up to 1 in 10 people) include:

- decreases in the number of white blood cells or red blood cells (seen in tests),
- dehydration, weight loss,
- sleeplessness (insomnia), depression,
- headache, sleepiness, dizziness, abnormal sensation in the skin (numbness or tingling sensation), taste changes,
- eye irritation, increased tears, eye redness (conjunctivitis)
- inflammation of the veins (thrombophlebitis),
- shortness of breath, nose bleeds, cough, runny nose,
- cold sores or other herpes infections,
- infections of the lungs or respiratory system (e.g. pneumonia or bronchitis),
- bleeding from the gut, constipation, pain in upper abdomen, indigestion, excess wind, dry mouth
- skin rash, hair loss (alopecia), skin reddening, dry skin, itching (pruritus), skin discolouration, skin loss, skin inflammation, nail disorder
- pain in the joints, or in the limbs (extremities), chest or back,
- fever, swelling in the limbs, feeling ill
- problems with liver function (seen in blood tests) and increased blood bilirubin (excreted by the liver)

Uncommon side effects (may affect up to 1 in 100 people) include:

- blood infection, urinary tract infection, infection of the skin, infections in the nose and throat, fungal infections (including those of the mouth), influenza, gastroenteritis, tooth abscess,
- lumps under the skin (lipoma),
- decreases in blood cells including platelets, thinning of blood (seen in tests)
- allergy

- diabetes, decrease in blood potassium, malnutrition, increased blood triglycerides,
- confusional state, panic attacks, depressed mood, decreased libido,
- difficulty speaking, impaired memory, loss of movement coordination, balance disorder, fainting, nerve damage (neuropathy) and problems with sensation
- blurred or double vision,
- vertigo, ear pain.
- irregular heartbeat and palpitations (arrhythmias), chest pain and heart attack (infarction),
- blood clots in the deep veins, high or low blood pressure, hot flushes, cold limbs (extremities), purple spots on the skin
- blood clots in the veins in the lung (pulmonary embolism), collapsed lung, coughing up blood, asthma, shortness of breath on exertion,
- bowel obstruction, collection of fluid in the abdomen, inflammation of the small or large intestine, the stomach or the oesophagus, pain in the lower abdomen, abdominal discomfort, heartburn (reflux of food from the stomach), blood in the stool,
- jaundice (yellowing of skin and eyes)
- skin ulcer and blister, reaction of the skin with sunlight, reddening of palms, swelling or pain of the face
- joint swelling or stiffness, bone pain, muscle weakness or stiffness,
- fluid collection in the kidneys, increased frequency of urination during the night, incontinence, blood in the urine, increase in blood creatinine (sign of kidney dysfunction)
- unusual bleeding from the vagina
- swelling (oedema), chills and rigors

Rare side effects (may affect up to 1 in 1,000 people) include

- angioedema (swelling mainly of the face, lip, tongue or throat, itching and rashes)

Some of these side effects are more common when capecitabine is used with other medicines for the treatment of cancer. Other side-effects seen in this setting are the following:

Common side effects (may affect up to 1 in 10 people) include:

- decrease in blood sodium, magnesium or calcium, increase in blood sugar,
- nerve pain,
- ringing or buzzing in the ears (tinnitus), loss of hearing,
- vein inflammation,
- hiccups, change in voice,
- pain or altered/abnormal sensation in the mouth, pain in the jaw,
- sweating, night sweats,
- muscle spasm,
- difficulty in urination, blood or protein in the urine,
- bruising or reaction at the injection site (caused by medicines given by injection at the same time)

Rare side effects (may affect up to 1 in 1,000 people) include:

- narrowing or blockage of tear duct (lacrimal duct stenosis),
- liver failure,
- inflammation leading to dysfunction or obstruction in bile secretion (cholestatic hepatitis),
- specific changes in the electrocardiogram (QT prolongation),
- certain types of arrhythmia (including ventricular fibrillation, torsade de pointes, and bradycardia).
- eye inflammation causing eye pain and possibly eyesight problems
- inflammation of the skin causing red scaly patches due to an immune system illness

Very rare side effects (may affect up to 1 in 10,000 people) include:

- severe skin reaction such as skin rash, ulceration and blistering which may involve ulcers of the mouth, nose, genitalia, hands, feet and eyes (red and swollen eyes).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V*](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [\[Nationally completed name\]](#)

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Do not store above 30°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [\[Nationally completed name\]](#) contains

The active substance is capecitabine.

Each Capecitabine 150 mg film-coated tablet contains 150 mg of capecitabine.

Each Capecitabine 500 mg film-coated tablet contains 500 mg of capecitabine.

The other ingredients are:

- Tablet core: lactose monohydrate, hypromellose, microcrystalline cellulose, croscarmellose sodium, magnesium stearate
- Tablet film-coating: hypromellose, talc, titanium dioxide (E171), iron oxide red (E172)

What [\[Nationally completed name\]](#) looks like and contents of the pack

[\[Nationally completed name\]](#) 150 mg film-coated tablets:

Light pink film-coated tablet of modified oval shape (5.5 x 11.0 mm) with the marking “150” on one side.

[\[Nationally completed name 500 mg film-coated tablets\]:](#)

Pink film-coated tablet of modified oval shape (8.4 x 16.0 mm) with the marking “500” on one side.

[\[NL/H/2458/001-002\]](#)

[\[Nationally completed name\]](#) are packaged in PVC/PVDC– Al foil blisters or alternatively in Al-Al blisters placed into cardboard boxes containing 28, 30, 50, 56, 60, 80, 84, 90, 100, 110, 112, 120 and 180 film-coated tablets.

Not all pack sizes may be marketed.

[NL/H/2459/001-002]

[Nationally completed name] are packaged in PVC/PVDC– Al foil blisters or alternatively in Al-Al blisters placed into cardboard boxes containing 30, 50, 60, 70, 90, 100 and 120 film-coated tablets. Not all pack sizes may be marketed.

Houder van de vergunning voor het in de handel brengen en fabrikant

Houder van de vergunning voor het in de handel brengen

Sandoz B.V., Veluwezoom 22, 1327 AH Almere, Nederland

Fabrikant

Salutas Pharma GmbH
Otto-von-Guericke Allee 1
39179 Barleben
Duitsland

In het register ingeschreven onder:

Capecitabine Sandoz 150 mg - RVG 110880

Capecitabine Sandoz 500 mg - RVG 110881

Dit geneesmiddel is geregistreerd in lidstaten van de EEA onder de volgende namen:

Nederland:	Capecitabine Sandoz 150 mg, filmomhulde tabletten Capecitabine Sandoz 500 mg, filmomhulde tabletten
Oostenrijk:	Capecitabin Sandoz 150 mg - Filmtabletten Capecitabin Sandoz 500 mg - Filmtabletten
Cyprus:	Capecitabin Sandoz
Denemarken:	Capecitabine Sandoz
Estland:	Capecitabine Sandoz
Griekenland:	Capecitabin/ Sandoz
Frankrijk:	Capecitabine Sandoz 150 mg, comprimé pelliculé Capecitabine Sandoz 500 mg, comprimé pelliculé
Hongarije:	Capecitabin Sandoz 150 mg filmtabletta Capecitabin Sandoz 500 mg filmtabletta
Ierland:	Capecitabine Sandoz 150 mg Film-Coated Tablets Capecitabine Sandoz 500 mg Film-Coated Tablets
Letland:	Capecitabine Sandoz 150 mg apvalkotās tabletes Capecitabine Sandoz 500 mg apvalkotās tabletes
Malta:	Capecitabine Sandoz 150 mg Filmcoated Tablets Capecitabine Sandoz 500 mg Filmcoated Tablets
Zweden:	Capecitabine Sandoz
Slovenië:	Kapecitabin Sandoz 150 mg filmsko obložene tablete Kapecitabin Sandoz 500 mg filmsko obložene tablete
Verenigd Koninkrijk:	Capecitabine Sandoz 150 mg Filmcoated Tablets Capecitabine Sandoz 500 mg Filmcoated Tablets

Deze bijsluiter is voor het laatst goedgekeurd in februari 2021