

PACKAGE LEAFLET

Package leaflet: Information for the user

Ibuprofen (als lysine) Mylan OTC 200 mg Film-coated tablets **Ibuprofen (als lysine) Mylan OTC 400 mg Film-coated tablets** Ibuprofen (as lysine)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days in the case of fever or after 4 days for the treatment of pain.

What is in this leaflet:

1. What Ibuprofen (als lysine) Mylan OTC is and what it is used for.
2. What you need to know before you take Ibuprofen (als lysine) Mylan OTC.
3. How to take Ibuprofen (als lysine) Mylan OTC.
4. Possible side effects.
5. How to store Ibuprofen (als lysine) Mylan OTC.
6. Contents of the pack and other information.

1. What Ibuprofen (als lysine) Mylan OTC is and what it is used for

Ibuprofen (als lysine) Mylan OTC contains the active substance ibuprofen. It belongs to a group of medicines known as non-steroidal anti-inflammatory drugs (NSAIDs). NSAIDs provide relief by changing the body's response to pain and high temperature.

Ibuprofen (als lysine) Mylan OTC tablets are used for the symptomatic treatment of mild to moderate pain, such as headache, dental pain, period pain and fever and pain as the result of a cold.

2. What do you need to know before you take Ibuprofen (als lysine) Mylan OTC

Do not take Ibuprofen (als lysine) Mylan OTC :

- if you are allergic to ibuprofen or any other ingredients of this medicine (listed in section 6)
- if you have had an allergic reaction such as asthma, difficulty in breathing, swelling of the face, tongue or throat nettle rash, itchy runny nose to acetylsalicylic acid (ASA) or other NSAIDs
- if you have (or have had two or more episodes of) a stomach ulcer or bleeding
- if you have had gastrointestinal perforation or bleeding when taking NSAIDs
- if you are suffering from cerebrovascular or other active bleeding
- if you are suffering from unclarified blood-formation disturbances
- if you have severe dehydration (caused by vomiting, diarrhoea or insufficient fluid intake)
- if you have severe liver, kidney or heart failure
- if you are in the last 3 months of pregnancy.

[200 mg only]

Ibuprofen (als lysine) Mylan OTC contraindicated in children under 6 years (less than 20 kg body weight).

[400 mg only]

Ibuprofen (als lysine) Mylan OTC is contraindicated in adolescents under 40 kg body weight or children under 12 years.

Warnings and precautions:

Talk to your doctor or pharmacist before taking Ibuprofen (als lysine) Mylan OTC:

- if you have recently had major surgery
- if you have or have had asthma or allergic disease as shortness of breath may occur
- if you suffer from hayfever, nasal polyps or chronic obstructive respiratory disorders as an increased risk of allergic reactions exists. The allergic reactions may present as asthma attacks (so-called analgesic asthma), Quincke's oedema or urticaria
- if you have heart problems
- if you have a history of gastrointestinal disease (such as ulcerative colitis, Crohn's disease)
- if you have systemic lupus erythematosus or mixed connective tissue disease (a disease affecting the skin, joints and kidneys)
- if you have certain hereditary blood formation disorders (e.g. acute intermittent porphyria)
- if you have a blood clotting disorder
- if you are taking other NSAIDs. The use with concomitant NSAIDs, including cyclo-oxygenase-2 specific inhibitors, increases the risk of adverse reactions (see section "Other medicines and Ibuprofen (als lysine) Mylan OTC" below) and should be avoided
- if you have chicken pox (varicella) it is advisable to avoid use of Ibuprofen (als lysine) Mylan OTC
- if you are elderly
- if you have an infection - please see heading "Infections" below.

Patients with kidney or liver problems should first consult a doctor before taking ibuprofen tablets.

Anti-inflammatory/pain-killer medicines like ibuprofen may be associated with a small increased risk of heart attack ("myocardial infarction") or stroke, particularly when used at high doses. Do not exceed the recommended dose or duration of treatment (see section 3).

You should discuss your treatment with your doctor or pharmacist before taking Ibuprofen (als lysine) Mylan OTC if you:

- have heart problems including heart failure, angina (chest pain), or if you have had a heart attack, bypass surgery, peripheral artery disease (poor circulation in the legs or feet due to narrow or blocked arteries), or any kind of stroke (including 'mini-stroke' or transient ischaemic attack, 'TIA')
- have high blood pressure, diabetes, high cholesterol, have a family history of heart disease or stroke, or if you are a smoker.

Skin reactions

Serious skin reactions have been reported in association with Ibuprofen (als lysine) Mylan OTC treatment. You should stop taking Ibuprofen (als lysine) Mylan OTC and seek medical attention immediately, if you develop any skin rash, lesions of the mucous membranes, blisters or other signs of allergy since this can be the first signs of a very serious skin reaction. See section 4.

Infections

Ibuprofen (als lysine) Mylan OTC may hide signs of infections such as fever and pain. It is therefore possible that Ibuprofen (als lysine) Mylan OTC may delay appropriate treatment of infection, which may lead to an increased risk of complications. This has been observed in pneumonia caused by bacteria and bacterial skin infections related to chickenpox. If you take this medicine while you have an infection and your symptoms of the infection persist or worsen, consult a doctor without delay.

Undesirable effects may be minimised by using the minimum effective dose for the shortest period of time. The elderly are at increased risk of side effects.

In general the habitual use of (several sorts of) analgesics can lead to lasting severe kidney problems. This risk may be increased under physical strain associated with loss of salt and dehydration. Therefore it should be avoided. There is a risk of renal impairment in dehydrated children and adolescents.

Prolonged use of any type of painkiller for headaches can make them worse. If this situation is experienced or suspected, medical advice should be obtained and treatment should be discontinued. The diagnosis of medication overuse headache (MOH) should be suspected in patients who have frequent or daily headaches despite (or because of) the regular use of headache medications.

In prolonged administration of Ibuprofen (als lysine) Mylan OTC regular checking of your liver values, the kidney function, as well as the blood count, is required. Your doctor may ask you to have blood tests during treatment.

Other medicines and Ibuprofen (als lysine) Mylan OTC

What should you avoid when you are taking this medicine?

Ibuprofen (als lysine) Mylan OTC may affect or be affected by some other medicines. For example:

- medicines that are anti-coagulants (i.e. thin blood/prevent clotting e.g. aspirin/acetylsalicylic acid, warfarin, ticlopidine)
- medicines that reduce high blood pressure (ACE-inhibitors such as captopril, beta-blockers such as atenolol, angiotensin-II receptor antagonists such as losartan)

Some other medicines may also affect or be affected by treatment with Ibuprofen (als lysine) Mylan OTC. You should therefore always seek the advice of your doctor or pharmacist before you use Ibuprofen (als lysine) Mylan OTC with other medicines.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell them if you are taking:

- - Acetylsalicylic acid, or other NSAIDs (anti-inflammatories and analgesics) Since this may increase the risk of gastrointestinal ulcers or bleeding
 - Digoxin (for heart insufficiency) Since the effect of digoxin may be enhanced
 - Glucocorticoids (medicinal products containing cortisone or cortisone-like substances) Since this may increase the risk of gastrointestinal ulcers or bleeding
 - Anti-platelet agents Since this may increase the risk of bleeding
 - Acetylsalicylic acid (low dose) Since the blood-thinning effect may be impaired
 - Medicines for thinning the blood (such as warfarin) Since ibuprofen may enhance the effects of these medicines
 - Phenytoin (for epilepsy) Since the effect of phenytoin may be enhanced
 - Selective serotonin reuptake inhibitors (medicines used for depression) As these may increase the risk of gastrointestinal bleeding
 - Lithium (a medicine for manic depressive illness and depression) Since the effect of lithium may be enhanced
 - Probenecid and Sulfinpyrazones (medicines for treating gout) Since the excretion of ibuprofen may be delayed
 - Medicines for high blood pressure and water tablets Since ibuprofen may diminish the effects of these medicines and there could be a possible increased risk for the kidney

- Potassium sparing diuretics Since this may lead to hyperkalaemia (high potassium levels in the blood)
- Methotrexate (a medicine for cancer or rheumatism) Since the effect of methotrexate may be enhanced
- Tacrolimus and cyclosporin (immunosuppressive medicines) Since kidney damage may occur
- Zidovudine: (a medicine for treating HIV/AIDS) Since the use of ibuprofen may result in an increased risk of bleeding into a joint or a bleed that leads to swelling in HIV (+) haemophiliacs
- Sulfonylureas (antidiabetic medicines) Interactions may be possible
- Quinolone antibiotics Since the risk for convulsions may be increased
- Mifepristone (medicine prescribed for pregnancy termination) Since ibuprofen may reduce the effect of this medicine
- Bisphosphonates (medicines prescribed to treat osteoporosis) Since these may increase the risk of gastrointestinal ulcers or bleeding
- Oxpentifylline (pentoxifylline) (medicine prescribed to increase the blood flow to arms and legs) Since this may increase the risk of gastrointestinal bleeding
- Baclofen, a muscle relaxant Since the toxicity of baclofen may be enhanced

Low dose ASA

Do not take this medicine if you are taking ASA at doses of above 75 mg daily. If you are on low-dose ASA (up to 75 mg daily) speak to your doctor or pharmacist before you take Ibuprofen (als lysine) Mylan OTC.

Ibuprofen (als lysine) Mylan OTC with alcohol

Consumption of alcohol should be avoided while taking Ibuprofen (als lysine) Mylan OTC as it may intensify possible side effects.

Pregnancy, breast-feeding and fertility

Do not take Ibuprofen (als lysine) Mylan OTC in the last 3 months of pregnancy. as it could harm your unborn child or cause problems at delivery. It can cause kidney and heart problems in your unborn baby. It may affect your and your baby's tendency to bleed and cause labour to be later or longer than expected. You should not take <INVENTED NAME> during the first 6 months of pregnancy unless absolutely necessary and advised by your doctor. If you need treatment during this period or while you are trying to get pregnant, the lowest dose for the shortest time possible should be used. If taken for more than a few days from 20 weeks of pregnancy onward, <INVENTED NAME> can cause kidney problems in your unborn baby that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby. If you need treatment for longer than a few days, your doctor may recommend additional monitoring.

Ibuprofen (als lysine) Mylan OTC tablets belong to a group of medicines which may impair fertility in woman. This effect is reversible on stopping the medicine. It is unlikely that Ibuprofen (als lysine) Mylan OTC , used occasionally, will affect your chances of becoming pregnant, however tell your doctor before taking this medicine if you have problems becoming pregnant.

Driving and using machines

For short-term use and at normal dosage this medicine has no or negligible influence on the ability to drive and use machines. If side-effects such as tiredness, dizziness, drowsiness and visual disturbances occur, do not drive or operate machines. Alcohol consumption increases the risk of these side effects.

3. How to take Ibuprofen (als lysine) Mylan OTC

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The product is intended for short term use only. The lowest effective dose should be used for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen (see section 2).

You should not take Ibuprofen (als lysine) Mylan OTC for longer than 3 days in the case of headache or fever, or 4 days for the treatment of pain unless your doctor tells you to.

If symptoms persist or worsen consult your doctor.

[200 mg only]

Adults and adolescents over 40 kg body weight(12 years of age and above):

The recommended dose is 200 mg to 400 mg (1 or 2 tablets), up to three times a day as required. Leave at least four hours between 200 mg (1 tablet) doses and at least six hours between 400 mg (2 tablets) doses. Do not take more than 1200 mg (6 tablets) in 24 hours.

Use in children over 6 years (20 kg – 40 kg body weight):

The recommended daily dose is 20 mg to 30 mg of ibuprofen per kg body weight divided into three or four separate doses as shown in the table below. Leave at least six to eight hours between doses.

| Body weight | Single dose | Maximum daily dose |
|--------------------|---|--|
| 20 kg – 29 kg | one 200 mg tablet (equals 200 mg ibuprofen) | three 200 mg tablets (equals 600 mg ibuprofen) |
| 30 kg – 39 kg | one 200 mg tablet (equals 200 mg ibuprofen) | four 200 mg tablets (equals 800 mg ibuprofen) |

If in children aged from 6 years and in adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a doctor should be consulted.

Use in children under 6 years (less than 20 kg body weight):

Ibuprofen (als lysine) Mylan OTC is contraindicated for use in children under 6 years (less than 20 kg body weight).

[400 mg only]

Adults and adolescents from 40 kg body weight (12 years of age and above)

The recommended dose is 400 mg (1 tablet), up to three times a day as required. Leave at least six hours between 400 mg (1 tablet) doses. Do not take more than 1200 mg (3 tablets) in 24 hours.

If in children aged from 12 years and in adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a doctor should be consulted.

Use in adolescents under 40 kg body weight or in children under 12 years of age:

Ibuprofen (als lysine) Mylan OTC is contraindicated for use in adolescents under 40 kg body weight or in children under 12 years.

Method of administration

Ibuprofen (als lysine) Mylan OTC tablets are swallowed whole with plenty of water. **Do not** chew the tablets.

Patients with a sensitive stomach should take the tablets during a meal. Taking the tablets after a meal may delay the onset of pain or fever relief. Do not exceed the recommended dose or take more frequently than recommended.

If you take more Ibuprofen (als lysine) Mylan OTC than you should

If you have taken more X than you should, or if children have been taken medicine by accident always contact a doctor or nearest hospital to get an opinion of the risk and advice on action to be taken.

The symptoms can include nausea, stomach pain, vomiting (may be blood streaked), headache, ringing in the ears, confusion and shaky eye movement. At high doses, drowsiness, chest pain, palpitations, loss of consciousness, convulsions (mainly in children), weakness and dizziness, blood in urine, cold body feeling, and breathing problems have been reported.

If you forget to take Ibuprofen (als lysine) Mylan OTC

Simply refer to the directions above on how to take the medicine and **do not** take more than is advised.

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects may be minimised by taking the lowest dose for the shortest time necessary to relieve the symptoms. Elderly people using this product are at increased risk of developing problems associated with side effects.

Medicines such as Ibuprofen (als lysine) Mylan OTC may be associated with a small increased risk of heart attack (“myocardial infarction”) or stroke.

Some of the undermentioned undesirable effects are less frequent when the maximum daily dose is 1200 mg compared to high-dose therapy in rheumatism patients.

If you think you have any of the following side effects or symptoms, stop taking this medicine and seek immediate help:

- stomach and intestinal ulcers, sometimes with bleeding and perforation, vomiting blood or have black tar-like stools (common: may affect up to 1 in 10 people)
- kidney disease with blood in the urine which can be associated with renal failure (uncommon: may affect up to 1 in 100 people)
- severe allergic reactions (very rare: may affect up to 1 in 10,000 people) such as:
 - difficulties in breathing or unexplained wheezing
 - dizziness or faster heartbeat
 - drop in blood pressure leading to shock
 - swelling of your face, tongue or throat
- potentially life-threatening skin rashes with severe blisters and bleeding in the lips, eyes, mouth, nose and genitals (Steven-Johnson syndrome) or serious skin reactions which starts with painful red areas, then large blisters and ends with peeling of layers of skin. This is accompanied by fever and chills, aching muscles and generally feeling unwell (toxic epidermal necrolysis) (very rare: may affect up to 1 in 10,000 people)
- severe condition of the skin that may affect the mouth and other parts of the body with symptoms including: red, often itchy spots, similar to the rash of measles, which starts on the limbs and sometimes on the face and the rest of the body. The spots may blister or may progress to form raised,

- red, pale-centred marks. Those affected may have fever, sore throat, headache and/or diarrhoea (very rare: may affect up to 1 in 10,000 people)
- severe flaking or peeling of the skin (very rare: may affect up to 1 in 10,000 people)
 - a severe skin reaction known as DRESS syndrome can occur. Symptoms of DRESS include: skin rash, fever, swelling of lymph nodes and an increase of eosinophils (a type of white blood cells) (not known: frequency cannot be estimated from the available data).
 - inflammation of the pancreas with severe upper stomach pain, often with nausea and vomiting (very rare: may affect up to 1 in 10,000 people)
 - nausea, vomiting, loss of appetite, feeling generally unwell, fever, itching, yellowing of the skin and eyes, light coloured bowel motions, dark coloured urine which may be signs of hepatitis or liver failure (very rare: may affect up to 1 in 10,000 people)
 - disease of the heart with shortness of breath and swelling of the feet or legs due to fluid build-up (heart failure) (not known: frequency cannot be estimated from the available data)
 - aseptic meningitis (infection around the brain or spinal cord with symptoms including fever, nausea, vomiting, headache, stiff neck and extreme sensitivity to bright light and clouding of the mental state and thus not being fully in contact with the environment) (very rare: may affect up to 1 in 10,000 people)
 - heart attack (“myocardial infarction”, very rare: may affect up to 1 in 10,000 people) or stroke (not known: frequency cannot be estimated from the available data)
 - severe kidney damage (papillary necrosis), particularly in long term use (rare: may affect up to 1 in 1,000 people)
 - worsening of infection-related inflammations (e.g. development of flesh eating bacteria syndrome) particularly if using other NSAIDs (very rare: may affect up to 1 in 10,000 people)

Stop taking the medicine and tell your doctor if you experience:

Very common (may affect more than 1 in 10 people)

- indigestion or heartburn
- abdominal pain (pains in your stomach) or other abnormal stomach

Not known (frequency cannot be estimated from the available data):

- Visual disturbances
- A red, scaly widespread rash with bumps under the skin and blisters mainly localized on the skin folds, trunk, and upper extremities accompanied by fever at the initiation of treatment (acute generalised exanthematous pustulosis). Stop using Ibuprofen (als lysine) Mylan OTC if you develop these symptoms and seek medical attention immediately. See also section 2.

Tell your doctor if you experience any of the below side effects:

Very common (may affect more than 1 in 10 people)

- nausea, wind, diarrhoea, constipation, vomiting

Common (may affect up to 1 in 10 people):

- headache, sleepiness, dizziness, spinning sensation tiredness, agitation, inability to sleep, irritability
- stomach and intestinal ulcers, sometimes with bleeding and perforation; hidden blood loss which may lead to a condition in which there is a decreased number of red blood cells (symptoms include tiredness, headaches, being short of breath when exercising, dizziness and looking pale), black tar-like stools, vomiting blood, mouth ulcers and cold sores, inflammation of the colon (symptoms include diarrhoea, usually with blood and mucus, stomach pain, fever), worsening of inflammatory bowel disease, inflammation of bowel wall

Uncommon (may affect up to 1 in 100 people):

- hives, itching, unusual bleeding or bruising under the skin, skin rash, asthma attacks (sometimes with hypotension)
- runny or blocked nose, sneezing, facial pressure or pain, difficulty in breathing
- inflammation of the stomach (symptoms include pain, nausea, vomiting, vomiting blood, blood in the bowel motions)
- increased sensitivity of the skin to sun

- swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing, sometime with high blood pressure
- fluid build-up in the body's tissues especially in patients with high blood pressure or kidney problems

Rare (may affect up to 1 in 1,000 people)

- a disease affecting the skin, joints and kidneys (lupus erythematosus syndrome)
- depression, confusion, hallucinations, mental illness with strange or disturbing thoughts or moods
- buzzing, hissing, whistling, ringing or other persistent noise in the ears
- increase of blood urea nitrogen, serum transaminases and alkaline phosphatase, decrease in haemoglobin and haematocrit values, inhibition of platelet aggregation, prolonged bleeding time, decrease of serum calcium, increase in serum uric acid, all seen on a blood test
- loss of vision

Very rare (may affect up to 1 in 10,000 people):

- fast or irregular heart beats (palpitations)
- fluid build-up in the body's tissues
- inflammation of the food pipe, narrowing of intestines
- liver disease, liver damage (especially in long-term use), liver failure, yellowing of the skin and/or eyes, also called jaundice
- severe skin infections with soft tissue complications may occur if you have chicken pox
- build-up of fluid in the lungs, symptoms include breathlessness, which may be very severe and usually worsens on lying down.

Not known (frequency cannot be estimated from available data)

- worsening of an inflammatory disease that can affect any part of the gastrointestinal tract (symptoms include pain, fever, diarrhoea and loss of weight), worsening of inflammation of the colon (symptoms diarrhoea, usually with blood and mucus, stomach pain, fever)
- high blood pressure
- unusual hair loss or thinning
- menstrual period disorders

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the national reporting system listed in Appendix V* [to be completed nationally]. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ibuprofen (als lysine) Mylan OTC

Keep this medicine out of the sight and reach of children.

Do not use Ibuprofen (als lysine) Mylan OTC after the expiry date which is stated on the blister, carton or bottle after 'EXP'. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Bottles: Use within 60 days of opening. Once open, keep bottle tightly closed.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ibuprofen (als lysine) Mylan OTC contains:

The active substance is ibuprofen (as lysine).
Each 200 mg tablet contains 200 mg of ibuprofen (as lysine).
Each 400 mg tablet contains 400 mg of ibuprofen (as lysine).

The other ingredients of a tablet are cellulose, microcrystalline; silica, colloidal anhydrous; crospovidone; povidone; magnesium stearate; talc (E553b). The tablet coating contains polyvinyl alcohol, hydrolysed; titanium dioxide (E171); macrogol (E1521); talc (E553b). The printing ink contains shellac; iron oxide black (E172); ammonium hydroxide (E527).

What Ibuprofen (als lysine) Mylan OTC looks like and contents of the pack

200 mg strength comes as a white to off-white, film-coated, round, biconvex, bevelled edge tablet imprinted with “M” over “IL1” in black ink on one side of the tablet and blank on the other side.

400 mg strength comes as a white to off-white, film-coated, oval, biconvex, bevelled edge tablet imprinted with “M IL2” in black ink on one side of the tablet and blank on the other side.

Ibuprofen (als lysine) Mylan OTC is available in blister packs of 2, 4, 6, 10, 12, 16, 20, 24, 30, 36, 48, 50, 100, 200 tablets and bottles of 48, 100, 200 (dispensing pack) tablets. Each bottle also contains silica gel (desiccant) which helps to protect your tablets against moisture.

Not all pack sizes may be marketed.

Dit middel is ingeschreven in het register onder nummer:
RVG 111362- Ibuprofen (als lysine) Mylan OTC 200 mg, filmomhulde tabletten
RVG 111363- Ibuprofen (als lysine) Mylan OTC 400 mg, filmomhulde tabletten

Marketing Authorisation Holder

Registratiehouder:
Mylan Pharmaceuticals Ltd
Damastown Industrial Park
Mulhuddart
Dublin 15
DUBLIN
Ierland

Voor informatie en inlichtingen:
Mylan B.V.
Krijgsman 20
Amstelveen

Manufacturers

Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland.
Mylan Hungary Kft, H-2900 Komarom, Mylan utca 1, Hungary.

This medicinal product is authorised in the Member States of the EEA under the following names:

| | |
|-----------------------|--|
| Belgium | Ibuprofen Lysine Mylan Pharma 200 mg filmomhulde tabletten Ibuprofen Lysine Mylan Pharma 400 mg filmomhulde tabletten |
| Germany | Ibuprofen Mylan 400 mg Filmtabletten |
| Ireland | Ibuprofen 200 mg film-coated tablets Ibuprofen 400 mg film-coated tablets |
| Italy | BRUFEN ANALGESICO |
| Luxembourg | Ibuprofen Lysine Mylan Pharma 200 mg comprimés pelliculés Ibuprofen Lysine Mylan Pharma 400 mg comprimés pelliculés |
| Poland | Myibuprofen Myibuprofen Forte |
| The Netherlands (RMS) | Ibuprofen (als lysine) Mylan OTC 200 mg, filmomhulde tabletten Ibuprofen (als lysine) Mylan OTC 400 mg, filmomhulde tabletten |

Deze bijsluiter is voor het laatst goedgekeurd in maart 2023.