

Package leaflet: Information for the user

Ivabradine Chemo Ibérica 5 mg filmomhulde tabletten Ivabradine Chemo Ibérica 7,5 mg filmomhulde tabletten ivabradine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Ivabradine Chemo Ibérica is and what it is used for
2. What you need to know before you take Ivabradine Chemo Ibérica
3. How to take Ivabradine Chemo Ibérica
4. Possible side effects
5. How to store Ivabradine Chemo Ibérica
6. Contents of the pack and other information

1. What Ivabradine Chemo Ibérica is and what it is used for

Ivabradine Chemo Ibérica (ivabradine) is a heart medicine used to treat:

- Symptomatic stable angina pectoris (which causes chest pain) in adult patients whose heart rate is over or equal to 70 beats per minute. It is used in adult patients who do not tolerate or cannot take heart medicines called beta-blockers. It is also used in combination with beta-blockers in adult patients whose condition is not fully controlled with a beta-blocker.
- Chronic heart failure in adult patients whose heart rate is over or equal to 75 beats per minute. It is used in combination with standard therapy, including beta-blocker therapy or when beta-blockers are contraindicated or not tolerated.

About stable angina pectoris (usually referred to as “angina”):

Stable angina is a heart disease which happens when the heart does not receive enough oxygen. It usually appears between 40 and 50 years of age. The most common symptom of angina is chest pain or discomfort. Angina is more likely to happen when the heart beats faster in situations such as exercise, emotion, exposure to the cold or after eating. This increase in heart rate can cause the chest pain in people who suffer from angina.

About chronic heart failure:

Chronic heart failure is a heart disease which happens when your heart cannot pump enough blood to the rest of your body. The most common symptoms of heart failure are breathlessness, fatigue, tiredness and ankle swelling.

How does Ivabradine Chemo Ibérica work?

<invented name> mainly works by reducing the heart rate by a few beats per minute. This lowers the heart's need for oxygen especially in the situations when an angina attack is more likely to happen. In this way <invented name> helps to control and reduce the number of angina attacks.

Furthermore as elevated heart rate adversely affects the heart functioning and vital prognosis in patients with chronic heart failure, the specific heart rate lowering action of ivabradine helps to improve the heart functioning and vital prognosis in these patients.

2. What you need to know before you take Ivabradine Chemo Ibérica

Do not take Ivabradine Chemo Ibérica

- if you are allergic to ivabradine or any of the other ingredients of this medicine (listed in section 6);
- if your resting heart rate before treatment is too slow (below 70 beats per minute);
- if you are suffering from cardiogenic shock (a heart condition treated in hospital);
- if you suffer from a heart rhythm disorder;
- if you are having a heart attack;
- if you suffer from very low blood pressure;
- if you suffer from unstable angina (a severe form in which chest pain occurs very frequently and with or without exertion);
- if you have heart failure which has recently become worse;
- if your heart beat is exclusively imposed by your pacemaker;
- if you suffer from severe liver problems;
- if you are already taking medicines for the treatment of fungal infections (such as ketoconazole, itraconazole), macrolide antibiotics (such as josamycin, clarithromycin, telithromycin or erythromycin given orally), medicines to treat HIV infections (such as nelfinavir, ritonavir) or nefazodone (medicine to treat depression) or diltiazem, verapamil (used for high blood pressure or angina pectoris);
- if you are a woman able to have children and not using reliable contraception;
- if you are pregnant or trying to become pregnant;
- if you are breast-feeding.

Warnings and precautions

Talk to your doctor or pharmacist before taking Ivabradine Chemo Ibérica

- if you suffer from heart rhythm disorders (such as irregular heartbeat, palpitation, increase in chest pain) or sustained atrial fibrillation (a type of irregular heartbeat), or an abnormality of electrocardiogram (ECG) called 'long QT syndrome',
- if you have symptoms such as tiredness, dizziness or shortness of breath (this could mean that your heart is slowing down too much),
- if you suffer from symptoms of atrial fibrillation (pulse rate at rest unusually high (over 110 beats per minute) or irregular, without any apparent reason, making it difficult to measure),
- if you have had a recent stroke (cerebral attack),
- if you suffer from mild to moderate low blood pressure,
- if you suffer from uncontrolled blood pressure, especially after a change in your antihypertensive treatment,
- if you suffer from severe heart failure or heart failure with abnormality of ECG called 'bundle branch block',
- if you suffer from chronic eye retinal disease,
- if you suffer from moderate liver problems,
- if you suffer from severe renal problems.

If any of the above applies to you, talk straight away to your doctor before or while taking Ivabradine Chemo Ibérica Children

Ivabradine Chemo Ibérica is not intended for use in children and adolescents younger than 18 years.

Other medicines and Ivabradine Chemo Ibérica

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Make sure to tell your doctor if you are taking any of the following medicines, as a dose adjustment of Ivabradine Chemo Ibérica or monitoring should be required:

- fluconazole (an antifungal medicine)

- rifampicin (an antibiotic)
- barbiturates (for difficult sleeping or epilepsy)
- phenytoin (for epilepsy)
- *Hypericum perforatum* or St John's Wort (herbal treatment for depression)
- QT prolonging medicines to treat either heart rhythm disorders or other conditions :
 - quinidine, disopyramide, ibutilide, sotalol, amiodarone (to treat heart rhythm disorders)
 - bepridil (to treat angina pectoris)
 - certain types of medicines to treat anxiety, schizophrenia or other psychoses (such as pimozide, ziprasidone, sertindole)
 - anti-malarial medicines (such as mefloquine or halofantrine)
 - intravenous erythromycin (an antibiotic)
 - pentamidine (an antiparasitic medicine)
 - cisapride (against the gastro-oesophageal reflux)
- Some types of diuretics which may cause decrease in blood potassium level, such as furosemide, hydrochlorothiazide, indapamide (used to treat oedema, high blood pressure).

Ivabradine Chemo Ibérica with food and drink

Avoid grapefruit juice during treatment with <invented name>.

heeft opmaak toegepast: Engels (Verenigde Staten)

Pregnancy and breast-feeding

Do not take Ivabradine Chemo Ibérica if you are pregnant or are planning to have a baby (see “Do not take Ivabradine Chemo Ibérica”).

heeft opmaak toegepast: Engels (Verenigde Staten)

heeft opmaak toegepast: Engels (Verenigde Staten)

If you are pregnant and have taken Ivabradine Chemo Ibérica ,talk to your doctor.

heeft opmaak toegepast: Engels (Verenigde Staten)

Do not take Ivabradine Chemo Ibérica if you are able to become pregnant unless you use reliable contraceptive measures (see “Do not take Ivabradine Chemo Ibérica”).

heeft opmaak toegepast: Engels (Verenigde Staten)

Do not take Ivabradine Chemo Ibérica if you are breast-feeding (see “Do not take Ivabradine Chemo Ibérica”). Talk to your doctor if you are breast-feeding or intending to breast-feed as breastfeeding should be discontinued if you take Ivabradine Chemo Ibérica.

heeft opmaak toegepast: Engels (Verenigde Staten)

heeft opmaak toegepast: Engels (Verenigde Staten)

heeft opmaak toegepast: Engels (Verenigde Staten)

heeft opmaak toegepast: Engels (Verenigde Staten)

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Ivabradine Chemo Ibérica may cause temporary luminous visual phenomena (a temporary brightness in the field of vision, see “Possible side effects”). If this happens to you, be careful when driving or using machines at times when there could be sudden changes in light intensity, especially when driving at night.

heeft opmaak toegepast: Engels (Verenigde Staten)

Ivabradine Chemo Ibérica contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

heeft opmaak toegepast: Engels (Verenigde Staten)

3. How to take Ivabradine Chemo Ibérica

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Ivabradine Chemo Ibérica should be taken during

meals. The tablet can be divided into equal doses

If you are being treated for stable angina pectoris

The starting dose should not exceed one tablet of Ivabradine Chemo Ibérica 5 mg twice daily. If you still have angina symptoms and if you have tolerated the 5 mg twice daily dose well, the dose may be increased. The maintenance dose should not exceed 7.5 mg twice daily. Your doctor will prescribe the right dose for you. The usual dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of Ivabradine Chemo Ibérica 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

If you are being treated for chronic heart failure

The usual recommended starting dose is one tablet of Ivabradine Chemo Ibérica 5 mg twice daily increasing if necessary to one tablet of Ivabradine Chemo Ibérica 7.5 mg twice daily. Your doctor will decide the right dose for you. The usual dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of

Ivabradine Chemo Ibérica 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

If you take more Ivabradine Chemo Ibérica than you should:

A large dose of Ivabradine Chemo Ibérica could make you feel breathless or tired because your heart slows down too much. If this happens, contact your doctor immediately.

If you forget to take Ivabradine Chemo Ibérica:

If you forget to take a dose of Ivabradine Chemo Ibérica, take the next dose at the usual time. Do not take a double dose to make up for the forgotten dose.

If you stop taking Ivabradine Chemo Ibérica:

As the treatment for angina or chronic heart failure is usually life-long, you should discuss with your doctor before stopping this medicinal product.

If you think that the effect of Ivabradine Chemo Ibérica is too strong or too weak, talk to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The frequency of possible side effects listed below is defined using the following convention:

Very common:	may affect more than 1 in 10 people
Common:	may affect up to 1 in 10 people
Uncommon:	may affect up to 1 in 100 people
Rare:	may affect up to 1 in 1,000 people
Very rare:	may affect up to 1 in 10,000 people
Not known:	frequency cannot be estimated from the available data

The most common adverse reactions with this medicine are dose dependent and related to its mode of action:

Very common:

Luminous visual phenomena (brief moments of increased brightness, most often caused by sudden changes in light intensity). They can also be described as a halo, coloured flashes, image decomposition or multiple images. They generally occur within the first two months of treatment after which they may occur repeatedly and resolve during or after treatment

Common:

Modification in the heart functioning (the symptoms are a slowing down of the heart rate). They particularly occur within the first 2 to 3 months of treatment initiation.

Other side effects have also been reported:

Common:

Irregular rapid contraction of the heart, abnormal perception of heartbeat, uncontrolled blood pressure, headache, dizziness and blurred vision (cloudy vision).

Uncommon:

Palpitations and cardiac extra beats, feeling sick (nausea), constipation, diarrhoea, abdominal pain, spinning sensation (vertigo), difficulty breathing (dyspnoea), muscle spasms, changes in laboratory parameters : high blood levels of uric acid, an excess of eosinophils (a type of white blood cell) and elevated creatinine in blood (a breakdown product of muscle), skin rash, angioedema (such as swollen face, tongue or throat, difficulty in breathing or swallowing), low blood pressure, fainting, feeling of tiredness, feeling of weakness, abnormal ECG heart tracing, double vision, impaired vision.

Rare:

Urticaria, itching, skin reddening, feeling unwell.

Very rare:

Irregular heart beats.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ivabradine Chemo Ibérica

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicine via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Ivabradine Chemo Ibérica contains

- The active substance is ivabradine (as hydrochloride).

One film-coated tablet contains ivabradine hydrochloride equivalent to 5 mg of ivabradine

One film-coated tablet contains ivabradine hydrochloride equivalent to 7,5 mg of ivabradine

The other ingredients in the tablet core are: lactose monohydrate, magnesium stearate (E 470 B), maize starch, maltodextrin, colloidal anhydrous silica (E 551), and in the tablet coating: lactose monohydrate, hypromellose (E 464), titanium dioxide (E 171), macrogol (E-1521), yellow iron oxide (E 172), red iron oxide (E 172).

What Ivabradine Chemo Ibérica looks like and contents of the pack

Film-coated tablet.

Ivabradine Chemo Ibérica 5 mg film-coated tablets

Ivabradine Chemo Ibérica 5 mg is pale orange, capsule shape (8,4 x 3,4 mm), biconvex film-coated tablet scored in one side.

The tablet can be divided into equal doses.

Ivabradine Chemo Ibérica 7.5 mg film-coated tablets

Ivabradine Chemo Ibérica 7.5 mg is pale orange, round (7,1 mm x 3,8 mm), biconvex film-coated

tablet. The tablets are available in blisters containing packs of 14, 28, 56, 84, 98, 100 and 112 tablets.

Not all pack sizes may be marketed.

Houder van de vergunning voor het in de handel brengen en fabrikant

Vergunninghouder:

CHEMO IBÉRICA, S.A.
Gran Via Carlos III, 98, 7º,
08028 Barcelona (Spain)

Fabrikant:

Laboratorios Licons, S.A.
Avda. Miralcampo 7
Poligono Industrial Miralcampo.
19200 Azuqueca de Henares – Guadalajara
Spanje

In het register ingeschreven onder:

Ivabradine Chemo Ibérica 5 mg filmomhulde tabletten: RVG 118340
Ivabradine Chemo Ibérica 7,5 mg filmomhulde tabletten: RVG 118341

Dit geneesmiddel is geregistreerd in lidstaten van de EEA onder de volgende namen:

Nederland: Ivabradine Chemo Ibérica 5/7,5 mg filmomhulde tabletten
Spanje: Ivabradina Chemo Ibérica 5/7,5 mg comprimidos recubiertos con película EFG
Verenigd Koninkrijk: Ivabradine 5/7,5 mg film-coated tablets

Deze bijsluiter is voor het laatst goedgekeurd in augustus 2021