

## Package leaflet: Information for the user

### Ulipristal Aristo 30 mg filmomhulde tablet

Ulipristal acetate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

Always take this medicine exactly as described in this leaflet or as your pharmacist, doctor or other healthcare professional has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your pharmacist, doctor, or other healthcare professional. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Ulipristal Aristo is and what it is used for
2. What you need to know before you take Ulipristal Aristo
3. How to take Ulipristal Aristo
4. Possible side effects
5. How to store Ulipristal Aristo
6. Contents of the pack and other information
  - Useful information about contraception

#### **1. What Ulipristal Aristo is and what it is used for**

##### Ulipristal Aristo is an emergency contraceptive

Ulipristal Aristo is a contraceptive intended to prevent pregnancy after unprotected sex or if your contraceptive method has failed. For example:

- if you had sex without protection;
- if your or your partner's condom tore, slipped or came off, or if you forgot to use one;
- if you did not take your contraceptive pill as recommended.

You should take the tablet as soon as possible after sex, and within a maximum of 5 days (120 hours).

This is because it is more effective if you take it as soon as possible after unprotected sex.

This medicine is suitable for any woman of childbearing age, including adolescents.

You can take the tablet at any time in the menstrual cycle.

##### Ulipristal Aristo does not work if you are already pregnant.

If your menstrual period is late, there is a possibility that you may be pregnant. When your period is late or when you have symptoms of pregnancy (heavy breasts, morning sickness) you should consult a doctor or other healthcare professional before taking the tablet.

If you have unprotected sex after taking the tablet, it will not stop you from becoming pregnant. Unprotected sex at any time during your cycle can lead to pregnancy.

### Ulipristal Aristo is not to be used for regular contraception

If you do not have a regular method of contraception, talk to your doctor or healthcare professional to choose one that is suitable for you.

### How Ulipristal Aristo works

Ulipristal Aristo contains the substance *ulipristal acetate* which acts by modifying the activity of the natural hormone progesterone which is necessary for ovulation to occur. As a result, this medicine works by postponing ovulation. Emergency contraception is not effective in every case. Of 100 women who take this medicine approximately 2 will become pregnant.

This medicine is a contraceptive used to prevent a pregnancy from starting. If you are already pregnant, it will not interrupt an existing pregnancy.

### Emergency contraception does not protect against sexually transmitted infections

Only condoms can protect you from sexually transmitted infections. This medicine will not protect you against HIV infection or any other sexually transmitted diseases (e.g. chlamydia, genital herpes, genital warts, gonorrhoea, hepatitis B and syphilis). Ask a healthcare professional for advice if you are worried about this.

There is more information about contraception at the end of this leaflet.

## **2. What you need to know before you take Ulipristal Aristo**

### **Do not take Ulipristal Aristo**

- if you are allergic to ulipristal acetate or any of the other ingredients of this medicine (listed in section 6).

### **Warning and precautions**

Talk to your pharmacist, doctor or other healthcare professional before taking this medicine

- if your period is late or you have symptoms of pregnancy (heavy breasts, morning sickness), as you may already be pregnant (see section “Pregnancy, breast-feeding and fertility”);
- if you suffer from severe asthma;
- if you suffer from severe liver disease.

In all women, emergency contraception should be taken as soon as possible after unprotected intercourse. There is some evidence that this medicine may be less effective with increasing body weight or body mass index (BMI), but these data were limited and inconclusive. Therefore, Ulipristal Aristo is still recommended for all women regardless of their weight or BMI.

You are advised to speak to a healthcare professional if you are concerned about any problems related to taking emergency contraception.

If you become pregnant despite taking this medicine, it is important that you see your doctor. See section “Pregnancy, breast-feeding and fertility” for more information.

### **Other contraceptives and Ulipristal Aristo**

This medicine may make regular hormonal contraceptives, like pills and patches, temporarily less effective.

If you are currently taking hormonal contraception, continue to use it as usual after taking the tablet, but be sure to use condoms every time you have sex until your next period.

Do not take Ulipristal Aristo together with another emergency contraceptive pill that contains levonorgestrel.

By taking them both together, you might make this medicine less effective.

### **Other medicines and Ulipristal Aristo**

Tell your pharmacist, doctor or other healthcare professional if you are taking or have recently taken any other medicines, including medicines obtained without a prescription or herbal medicines.

Some medicines may prevent Ulipristal Aristo from working effectively. If you have used any of the medicines listed below during the last 4 weeks, Ulipristal Aristo may be less suitable for you. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD):

- medicines used to treat epilepsy (for example, primidone, phenobarbital, phenytoin, fosphenytoine, carbamazepine, oxcarbazepine and barbiturates)
- medicines used to treat tuberculosis (for example, rifampicin, rifabutin)
- a treatment for HIV (ritonavir, efavirenz, nevirapine)
- a medicine used to treat fungal infections (griseofulvin)
- herbal remedies containing St John's wort (*Hypericum perforatum*).

Speak to your doctor or pharmacist before using Ulipristal Aristo when you use (or have recently used) any of the medicines stated above.

### **Pregnancy, breast-feeding and fertility**

#### Pregnancy

Before taking this medicine, if your period is late, tell your pharmacist, doctor or other healthcare professional, or do a pregnancy test in order to make sure you are not already pregnant (see section “Warning and precautions”).

This medicine is a contraceptive used to prevent a pregnancy from starting. If you are already pregnant it will not interrupt an existing pregnancy.

If you become pregnant despite taking this medicine, there is no evidence that it will affect your pregnancy. However, it is important that you see your doctor. As for any pregnancy, your doctor may want to check that the pregnancy is not outside the womb. This is especially important if you have severe abdominal (stomach) pain or bleeding or if you have previously had a pregnancy outside the womb, tubal surgery or long term (chronic) genital infection.

If you become pregnant despite taking Ulipristal Aristo, you are encouraged to ask your doctor to register your pregnancy in an official registry. You can also report this information on your own at <http://www.ulipristal-pregnancy-registry.com>. Your information will remain anonymous – nobody will know it is information about you. Sharing your information may help women in the future understand the safety or risks of Ulipristal Aristo during a pregnancy.

#### Breast-feeding

If you take this medicine while you are breast-feeding a baby, do not breast-feed for one week after taking this medicine. During this time, it is recommended to use a breast pump in order to maintain milk production, but throw away your breast milk. The effect of breast-feeding your baby in the week after taking this medicine is not known.

### Fertility

This medicine will not affect your future fertility. If you have unprotected sex after taking this medicine, it will not stop you from becoming pregnant. Therefore it is important you use condoms until your next period.

If you wish to start or continue with a regular method of contraception after using this medicine, you can do so but you should also use condoms until your next period.

### **Driving and using machines**

After taking this medicine, some women experience dizziness, drowsiness, blurred vision and/or loss of concentration (see section 4 “Possible side effects”). If you experience these symptoms, do not drive or use machines.

### **Ulipristal Aristo contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Ulipristal Aristo contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially ‘sodium-free’.

## **3. How to take Ulipristal Aristo**

Always take this medicine exactly as described in this leaflet or as your pharmacist, doctor or other healthcare professional has told you. Check with your pharmacist or doctor if you are not sure.

### How to take the Ulipristal Aristo film-coated tablet

- Take one tablet by mouth as soon as possible and no later than 5 days (120 hours) after unprotected sex or contraceptive failure. Take the tablet without delay.
- You can take the tablet at any time in your cycle.
- You can take the tablet at any time of the day either before, during or after a meal.
- If you are using one of the medicines that may prevent Ulipristal Aristo from working properly (see section 2 “What you need to know before you take Ulipristal Aristo”) or if you have used one of these medicines in the past 4 weeks, Ulipristal Aristo may work less effectively for you. Speak to your doctor or pharmacist before using Ulipristal Aristo. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD).

### If you vomit after taking Ulipristal Aristo

If you vomit (be sick, throw up) within 3 hours of taking the tablet, take another tablet as soon as possible.

### If you have sex again after taking Ulipristal Aristo

If you have unprotected sex after taking the tablet, it will not stop you from becoming pregnant. After you take the tablet and until your next period comes, you should use condoms every time you have sex.

#### If your next period is late after taking Ulipristal Aristo

After taking the tablet, it is normal for your next period to be a few days late. However, if your period is more than 7 days late; if it is unusually light or unusually heavy; or if you experience symptoms such as abdominal (stomach) pain, breast tenderness, vomiting or nausea, you may be pregnant. You should do a pregnancy test right away. If you are pregnant, it is important that you see your doctor. (See section “Pregnancy, breast-feeding and fertility”)

#### If you take more Ulipristal Aristo than you should

There have been no reports of harmful effects from taking a higher dose than recommended of this medicine. However, do ask your pharmacist, doctor or other healthcare professional for advice. If you have any further questions on the use of this medicine, ask your pharmacist, doctor or other healthcare professional.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some symptoms such as breast tenderness and abdominal (stomach) pain, throwing up (vomiting), feeling sick (nausea) are also possible signs of pregnancy. If you miss your period and experience such symptoms after taking Ulipristal Aristo, you should do a pregnancy test (See section “Pregnancy, breastfeeding and fertility”).

#### Common side effects (may affect up to 1 in 10 people)

- nausea, abdominal (stomach) pain or discomfort, vomiting
- painful periods, pelvic pain, breast tenderness
- headache, dizziness, mood swings
- muscle pain, back pain, tiredness

#### Uncommon side effects (may affect up to 1 in 100 people)

- diarrhoea, heartburn, wind, dry mouth
- unusual or irregular vaginal bleeding, heavy/prolonged periods, premenstrual syndrome, vaginal irritation or discharge, lesser or greater sex drive
- hot flushes
- appetite changes, emotional disorders, anxiety, agitation, trouble sleeping, sleepiness, migraine, visual disturbances
- influenza
- acne, skin lesions, itching
- fever, chills, malaise

#### Rare side effects (may affect up to 1 in 1,000 people)

- genital pain or itching, pain during sex, rupture of an ovarian cyst, unusually light period
- loss of concentration, vertigo, shaking, disorientation, fainting
- unusual sensation in eye, red eye, sensitivity to light
- dry throat, disturbance in taste
- allergic reactions such as rash,
- hives (itchy rash), or swelling of the face
- feeling thirsty

#### Reporting of side effects

If you get any side effects, talk to your pharmacist, doctor or other healthcare professional. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can provide more information on the safety of this medicine.

### **5. How to store Ulipristal Aristo**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

Store in the original package in order to protect from light.

This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via waste water. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Ulipristal Aristo contains**

- The active substance is ulipristal acetate. Each film-coated tablet contains 30 milligrams of ulipristal acetate.
- The other ingredients are lactose monohydrate, pregelatinised starch (Maize), sodium starch glycolate, magnesium stearate, hypromellose (E464), hydroxypropylcellulose (E463), stearic acid (E570), talc (E553b), titanium dioxide (E171).

#### **What Ulipristal Aristo looks like and contents of the pack**

Ulipristal Aristo is a white, round, biconvex film-coated tablet of 9.0 - 9.2 mm diameter embossed with “U30” on one side.

Ulipristal Aristo is available in a carton containing one transparent PVC/PVDC/aluminium blister of 1 film-coated tablet.

#### **Marketing Authorisation Holder**

Aristo Pharma GmbH  
Wallenroder Str. 8-10  
13435 Berlin  
Duitsland

**Manufacturer**

Aristo Pharma GmbH  
Wallenroder Str. 8-10  
13435 Berlin  
Duitsland

Haupt Pharma Münster GmbH  
Schleebrüggenkamp 15  
48159 Münster  
Duitsland

**Marketing Authorisation Number(s)**

RVG 122046

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Germany:	Ulipristal Aristo 30 mg Filmtablette
Netherlands:	Ulipristal Aristo 30 mg filmomhulde tablet
Austria:	Ulipristal Aristo 30 mg Filmtablette
Czech Republic:	Ulipristal acetate Aristo 30 mg potahovaná tableta
Italy:	Ulipristal Aristo 30 mg Compressa rivestita con film
Poland:	Ulipristal Aristo 30 mg tabletko powlekana
United Kingdom:	Ulipristal Aristo 30 mg film-coated tablet
Spain:	Ulipristal Aristo 30 mg comprimido recubierto con película EFG
Portugal:	Ulipristal Aristo 30 mg Comprimido revestido por película

**Deze bijsluiter is voor het laatst goedgekeurd in maart 2022.**

## USEFUL INFORMATION ABOUT CONTRACEPTION

### MORE ABOUT EMERGENCY CONTRACEPTION

The sooner you take emergency contraception, the better the chance of avoiding pregnancy. Emergency contraception will not affect your fertility.

Emergency contraception can delay ovulation within a given menstrual cycle, but it will not stop you from becoming pregnant if you have unprotected sex again. After you take emergency contraception and until your next period comes, you should use a condom every time you have sex.

### MORE ABOUT REGULAR CONTRACEPTION

If you have taken emergency contraception and you do not use a regular contraceptive method (or do not have a contraceptive method that suits you), talk to your doctor or family planning clinic for advice.

There are many different types of contraception available, and you should be able to find the right method for you.

Examples of regular contraception methods:

---

#### Daily methods

Contraceptive pill

---

#### Weekly or monthly methods

Contraceptive patch

Vaginal Ring

---

#### Long lasting methods

Contraceptive implant

IUD (intrauterine device)

---