

Package leaflet: Information for the patient

Quator[®] 2,5 mg, filmomhulde tabletten
Quator[®] 5 mg, filmomhulde tabletten
Quator[®] 10 mg, filmomhulde tabletten
Quator[®] 20 mg, filmomhulde tabletten

tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Nationally completed name] is and what it is used for
2. What you need to know before you take [Nationally completed name]
3. How to take [Nationally completed name]
4. Possible side effects
5. How to store [Nationally completed name]
6. Contents of the pack and other information

1. What [Nationally completed name] is and what it is used for

[Nationally completed name] contains the active substance tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors.

[Nationally completed name] **2.5 mg, 5 mg, 10 mg and 20 mg** film-coated tablets is used to treat adult men with **erectile dysfunction**. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. [Nationally completed name] has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.

Following sexual stimulation [nationally completed name] works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. [Nationally completed name] will not help you if you do not have erectile dysfunction. It is important to note that [Nationally completed name] for the treatment of erectile dysfunction does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

2. What you need to know before you take [Nationally completed name]

Do not take [Nationally completed name] if you:

- are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6)

- are taking any form of organic nitrate or nitric oxide donors such as amyl nitrite. This is a group of medicines (“nitrates”) used in the treatment of angina pectoris (“chest pain”). Tadalafil has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor.
- have serious heart disease or recently had a heart attack within the last 90 days
- recently had a stroke within the last 6 months
- have low blood pressure or uncontrolled high blood pressure
- ever had loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as “stroke of the eye”
- are taking riociguat. This medicine is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as [Nationally completed name], have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor before taking [Nationally completed name].

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Before taking the tablets, tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells)
- multiple myeloma (cancer of the bone marrow)
- leukaemia (cancer of the blood cells)
- any deformation of your penis
- a serious liver problem
- a severe kidney problem

It is not known if [Nationally completed name] is effective in patients who have had:

- pelvic surgery
- removal of all or part of the prostate gland in which nerves of the prostate are cut (radical non-nerve-sparing prostatectomy)

If you experience sudden decrease or loss of vision, or your vision is distorted, dimmed while you are taking [Nationally completed name] stop taking [Nationally completed name] and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking [Nationally completed name] and contact your doctor immediately.

[Nationally completed name] is not intended for use by women.

Children and adolescents

[Nationally completed name] is not intended for use by children and adolescents under the age of 18.

Other medicines and [Nationally completed name]

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do not take [Nationally completed name] if you are already taking nitrates.

Some medicines may be affected by [Nationally completed name] or they may affect how well [Nationally completed name] will work. Tell your doctor or pharmacist if you are already taking:

- an alpha-blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia)
- other medicines to treat high blood pressure
- riociguat
- a 5- alpha reductase inhibitor (used to treat benign prostatic hyperplasia)
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines)
- rifampicin, erythromycin , clarithromycin or itraconazole
- other treatments for erectile dysfunction

[Nationally completed name] with drink and alcohol

Information on the effect of alcohol is in section 3. Grapefruit juice may affect how well [Nationally completed name] will work and should be taken with caution. Talk to your doctor for further information.

Fertility

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Some men taking [Nationally completed name] in clinical studies have reported dizziness. Check carefully how you react to the tablets before driving or using machines.

[Nationally completed name] contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

[Nationally completed name] contains sodium This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take [Nationally completed name]

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Nationally completed name] tablets are for oral use in men only. Swallow the tablet whole with some water. [Nationally completed name] 20 mg tablets can be divided into equal doses. The tablets can be taken with or without food.

- **Flexible-dose, on demand, using 10 mg or 20 mg film-coated tablets:**

The recommended starting dose is one 10 mg tablet before sexual activity. If the effect of this dose is too weak your doctor may increase the dose to 20 mg. [Nationally completed name] tablets are for oral use.

You may take a [Nationally completed name] tablet at least 30 minutes before sexual activity. [Nationally completed name] may still be effective up to 36 hours after taking the tablet. [Nationally completed name] 10 mg and 20 mg is intended for use prior to anticipated sexual activity and is not recommended for continuous daily use.

- **Once a day dosing, using 2.5 mg or 5 mg film-coated tablets:**

The recommended dose is one 5 mg tablet taken once a day at approximately the same time of the day. Your doctor may adjust the dose to 2.5 mg based on your response to [Nationally completed name]. This will be given as a 2.5 mg tablet.

When taken once a day [Nationally completed name] allows you to obtain an erection, when sexually stimulated, at any time point during the 24 hours of the day. Once a day dosing of [Nationally completed name] may be useful to men who anticipate having sexual activity two or more times per week.

Do not take [Nationally completed name] more than once a day.

It is important to note that [Nationally completed name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Drinking alcohol may affect your ability to get an erection and may temporarily lower your blood pressure. If you have taken or are planning to take [Nationally completed name], avoid excessive drinking (blood alcohol level of 0.08% or greater), since this may increase the risk of dizziness when standing up.

If you take more [Nationally completed name] than you should

Contact your doctor. You may experience side effects described in section 4.

If you forget to take [Nationally completed name]

If you have a once a day dosing of [nationally completed name], take your dose as soon as you remember but do not take a double dose to make up for a forgotten dose. You should not take [Nationally completed name] more than once a day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects stop using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon, may affect up to 1 in 100 people)
- chest pain - do not use nitrates but seek immediate medical assistance (frequency uncommon, may affect up to 1 in 100 people)

- priapism, a prolonged and possibly painful erection after taking tadalafil (frequency rare, may affect up to 1 in 1,000 people)
- If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare, may affect up to 1 in 1,000 people), distorted, dimmed, blurred central vision or sudden decrease of vision (frequency not known)

Other side effects have been reported:

Common (may affect up to 1 in 10 people)

- headache, back pain, muscle aches, pain in arms and legs, facial flushing, nasal congestion and indigestion

Uncommon (may affect up to 1 in 100 people)

- dizziness, stomach ache, feeling sick, being sick (vomiting), reflux, blurred vision, eye pain, difficulty in breathing, presence of blood in urine, prolonged erection, pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure, nose bleeds, ringing in the ears, swelling of the hands, feet or ankles and feeling tired

Rare (may affect up to 1 in 1,000 people)

- fainting, seizures and passing memory loss, swelling of the eyelids, red eyes, sudden decrease or loss of hearing, hives (itchy red welts on the surface of the skin), penile bleeding, presence of blood in semen and increased sweating

Heart attack and stroke have also been reported rarely in men taking tadalafil. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking tadalafil that were not seen in clinical studies. These include:

- migraine, swelling of the face, serious allergic reaction which causes swelling of the face or throat, serious skin rashes, some disorders affecting blood flow to the eyes, irregular heartbeats, angina and sudden cardiac death
- distorted, dimmed, blurred central vision or sudden decrease of vision (frequency not known).

The side effect dizziness has been reported more frequently in men over 75 years of age taking tadalafil. Diarrhoea has been reported more frequently in men over 65 years of age taking tadalafil.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly **via the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [nationally completed name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [nationally completed name] contains

- The active substance is tadalafil.

<{[Nationally completed name] 2.5 mg film-coated tablets}>

Each film-coated tablet contains 2.5 mg of tadalafil.

<{[Nationally completed name] 5 mg film-coated tablets}>

Each film-coated tablet contains 5 mg of tadalafil.

<{[Nationally completed name] 10 mg film-coated tablets}>

Each film-coated tablet contains 10 mg of tadalafil.

<{[Nationally completed name] 20 mg film-coated tablets}>

Each film-coated tablet contains 20 mg of tadalafil.

- The other excipients are:

Tablet core:

Lactose monohydrate, sodium laurilsulfate (E487), povidone K-12 (E1201), crospovidone (Type B) (E1202), sodium stearyl fumarate. See section 2 [Nationally completed name] contains lactose.

Film-coating:

Polyvinyl alcohol (E1203), macrogol 3350 (E1521), titanium dioxide (E171), talc (E553b), iron oxide yellow (E172)

What [nationally completed name] looks like and contents of the pack

{[Nationally completed name] 2.5 mg film-coated tablets}

Ochre to yellow, round shaped film-coated tablet of approximately 5 mm in diameter. On one side debossed with "2.5" and plain on the other side.

{[Nationally completed name] 5 mg film-coated tablets}

Ochre to yellow, oval shaped film-coated tablet of approximately 8 x 4 mm. On one side debossed with "5" and plain on the other side.

{[Nationally completed name] 10 mg film-coated tablets}

Ochre to yellow, oval shaped film-coated tablet of approximately 11 x 6 mm. On one side debossed with "10". The other side of the tablet is plain.

{[Nationally completed name] 20 mg film-coated tablets}

Ochre to yellow, oval shaped film-coated tablet of approximately 15 x 9 mm. On one side debossed with "20". The other side of the tablet is bi-scored. The tablet can be divided into equal halves and/or quarters.

Alu- Alu (Al-OPA/Al/PVC) blisters
PVC/ACLAR/PVC - Alu blisters
PVC/ACLAR/PVdC/PVC KPMAX - Alu blisters

Pack sizes:

[NL/H/4395/001-004/DC]

2.5 mg: 28x1 tablets
5 mg: 14x1, 28x1, 84x1 tablets
10 mg: 4x1, 12x1, 24x1, 36x1, 48x1 tablets
20 mg: 2x1, 4x1, 8x1, 12x1, 14x1, 24x1, 28x1, 36x1, 48x1 tablets

[NL/H/4396/001-004/DC]

2.5 mg: 28x1 tablets
5 mg: 14x1, 28x1 tablets
10 mg: 4x1, 12x1, 24x1, 36x1, 48x1 tablets
20 mg: 2x1, 4x1, 8x1, 12x1, 14x1, 24x1, 28x1, 36x1, 48x1 tablets

[NL/H/4397/001-004/DC]

2.5 mg: 28x1 tablets
5 mg: 14x1, 28x1 tablets
10 mg: 4x1, 12x1, 24x1, 36x1, 48x1 tablets
20 mg: 2x1, 4x1, 8x1, 12x1, 14x1, 24x1, 28x1, 36x1, 48x1 tablets

[NL/H/4398/001-003/DC]

5 mg: 14x1 tablets
10 mg: 4x1 tablets
20 mg: 2x1, 4x1 tablets

Not all pack sizes may be marketed.

Houder van de vergunning voor het in de handel brengen en fabrikant

Houder van de vergunning voor het in de handel brengen

Sandoz B.V.
Veluwezoom 22
1327 AH Almere
Nederland

Fabrikant

TEVA PHARMA, S.L.U.
Polígono Industrial Malpica, C/C, nº4
50016, Zaragoza
Spanje

LEK PHARMACEUTICALS D.D.,
Verovškova ulica 57
Ljubljana
Slovenië

In het register ingeschreven onder:

Quator 2,5 mg, filmomhulde tabletten – RVG 123026
Quator 5 mg, filmomhulde tabletten – RVG 123027
Quator 10 mg, filmomhulde tabletten – RVG 123028
Quator 20 mg, filmomhulde tabletten – RVG 123029

Dit middel is geregistreerd in lidstaten van de EEA onder de volgende namen:

België Quator 2,5 mg filmomhulde tabletten
Quator 5 mg filmomhulde tabletten
Quator 10 mg filmomhulde tabletten
Quator 20 mg filmomhulde tabletten
Denemarken Quator
Duitsland Quator 2,5 mg Filmtabletten
Quator 5 mg Filmtabletten
Quator 10 mg Filmtabletten
Quator 20 mg Filmtabletten
Estland Quator
Frankrijk Quator 2,5 mg, comprimé pelliculé
Quator 5 mg, comprimé pelliculé
Quator 10 mg, comprimé pelliculé
Quator 20 mg, comprimé pelliculé
Griekenland Quator 5 mg επικαλυμμένα με λεπτό υμένιο δισκία
Quator 10 mg επικαλυμμένα με λεπτό υμένιο δισκία
Quator 20 mg επικαλυμμένα με λεπτό υμένιο δισκία
Ierland Quator 10 mg Film-coated tablets
Quator 20 mg Film-coated tablets
Letland Quator 5 mg apvalkotās tabletes
Quator 10 mg apvalkotās tabletes
Quator 20 mg apvalkotās tabletes
Litouwen Quator 5 mg plėvele dengtosd tabletės
Quator 10 mg plėvele dengtosd tabletės
Quator 20 mg plėvele dengtosd tabletės
Nederland Quator 2,5 mg, filmomhulde tabletten
Quator 5 mg, filmomhulde tabletten
Quator 10 mg, filmomhulde tabletten
Quator 20 mg, filmomhulde tabletten
Oostenrijk Quator 2,5 mg – Filmtabletten
Quator 5 mg – Filmtabletten
Quator 10 mg – Filmtabletten
Quator 20 mg – Filmtabletten
Portugal Quator

Slovenië Quator 5 mg filmsko obložene tablete
Quator 10 mg filmsko obložene tablete
Quator 20 mg filmsko obložene tablete
Slowakije Quator 5 mg
Quator 10 mg

Deze bijsluiter is voor het laatst goedgekeurd in april 2024