

Package leaflet: Information for the user

Lisdexamfetamine Teva 30 mg, harde capsules
Lisdexamfetamine Teva 50 mg, harde capsules
Lisdexamfetamine Teva 70 mg, harde capsules
lisdexamfetamine dimesylate

Important things you need to know about your medicine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4, for more information.

What is in this leaflet

1. What [Product name] is and what it is used for
2. What you need to know before you take [Product name]
3. How to take [Product name]
4. Possible side effects
5. How to store [Product name]
6. Contents of the pack and other information

1. What [Product name] is and what it is used for

What [Product name] is

[Product name] contains the active substance lisdexamfetamine dimesylate which helps with your brain activity. It helps improve your attention, helps you concentrate and makes you less impulsive. [Product name] is a long acting medicine which works gradually over a 14 hour time period.

What it is used for

[Product name] is a treatment for 'attention deficit hyperactivity disorder' (ADHD) in adults.

You must talk to a doctor if you do not feel better or if you feel worse after one month of treatment. For children and adolescents aged 6 to 17 years, another product containing lisdexamfetamine dimesylate is available. [Product name] is not used as a treatment for ADHD in children under 6 years of age because it is not known if it is safe or of benefit in such young people.

How it works

[Product name] improves the activity of certain parts of the brain which are under-active. The medicine can help improve attention, concentration and reduce impulsive behaviour.

The medicine is given as part of a treatment programme, which usually includes the following:

- psychological therapy
- educational therapy
- behavioural therapy
- occupational therapy

- social therapy

It is prescribed only by doctors who have experience in treating people with behaviour problems.

If you have not been treated for ADHD before, the doctor will check if you have had ADHD since childhood before prescribing [Product name].

About ADHD

People with ADHD find it hard to:

- sit still
- concentrate

It is not their fault that they cannot do these things. However, ADHD can cause problems with everyday life. People with ADHD may have difficulty working, learning, with relationships and have low self esteem.

ADHD does not affect the intelligence of a person

2. What you need to know before you take [Product name]

Do NOT take [Product name]:

- if you are allergic to lisdexamfetamine or any of the other ingredients of this medicine (listed in Section 6)
- if you are taking a medicine called a ‘monoamine oxidase inhibitor’ (MAOI) used for depression, or have taken an MAOI in the last 14 days
- if you have a thyroid problem
- if you feel unusually excited, over-active, or un-inhibited
- if you have ever had heart problems - such as a heart attack, uneven heartbeat, pain and discomfort in the chest, heart failure, heart disease or were born with a heart problem
- if you have high or very high blood pressure or narrowing of the blood vessels
- if you have increased pressure in your eye (glaucoma)

Do not take [Product name] if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before you take [Product name]. This is because [Product name] can make these problems worse.

Warnings and precautions

Talk to your doctor or pharmacist before using [Product name] treatment if you have:

- ever abused prescription medicines or street drugs
- had kidney problems
- had fits (seizures, convulsions, epilepsy) or any abnormal brain scans (EEGs)
- plan to become pregnant or are pregnant (see the ‘Pregnancy, breast-feeding and fertility’ section)
- hard-to-control and repeated twitching of any parts of the body or you repeat sounds and words
- high blood pressure
- family or medical history of irregular heart rhythm (visible on an electrocardiogram), or if you have a disease and/or take a treatment that make(s) you prone to heartbeat irregularities or salt imbalances
- a heart problem which is not in the ‘Do not take’ section above
- a mental health problem. These may include:
 - mood swings (from being manic to being depressed - called ‘bipolar disorder’)
 - starting to be aggressive or unfriendly (hostile), or your aggression gets worse
 - seeing, hearing or feeling things that are not there (hallucinations)
 - believing things that are not true (delusions)
 - feeling unusually suspicious (paranoia)

- feeling agitated, anxious or tense
- feeling depressed or guilty

Tell your doctor or pharmacist if any of the above applies to you before starting treatment. This is because [Product name] can make these problems worse. Your doctor will want to monitor how the medicine affects you.

[Product name] may cause heart rhythm disorders in some patients. If you experience palpitations or irregular heartbeat during the period of treatment, you should inform your doctor immediately. The risk of heart problems may increase with increase of the dose. Therefore, the recommended dosage should be followed.

Checks that your doctor will make before you start taking [Product name]

These checks are to decide if [Product name] is the correct medicine for you. Your doctor will ask you about:

- any other medicines you are taking
- whether there is any family history of sudden unexplained death
- any other medical problems (such as heart problems) you or your family may have
- how you are feeling, such as feeling happy or sad, having strange thoughts, or if you have had any of these feelings in the past
- whether there is a family history of ‘tics’ (hard-to-control, repeated twitching of any parts of the body or repeating sounds and words)
- any mental health or behaviour problems you or other family members have ever had. Your doctor will check your mental health history, and check if any of your family have a history of suicide, bipolar disorder (mood swings from being manic to being depressed) or depression.

It is important that you provide as much information as you can. This will help your doctor decide if [Product name] is the correct medicine for you. Your doctor may decide that other medical tests are needed before you start taking this medicine.

Other medicines and [Product name]

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

Do NOT take [Product name] if you:

- are taking a medicine called a ‘monoamine oxidase inhibitor’ (MAOI) used for depression, or have taken an MAOI in the last 14 days. Taking an MAOI with [Product name] may cause a sudden increase in your blood pressure. Your doctor or pharmacist will be able to tell you if are taking a medicine that is an MAOI.

[Product name] and some other medicines can affect each other. If you are taking any of the following medicines, check with your doctor or pharmacist before taking [Product name]:

- medicines for severe mental health problems
- medicines used to reduce or increase blood pressure
- medicines used during surgery such as painkillers
- cough and cold remedies. Some of these contain medicines that can affect blood pressure so it is important to check with your pharmacist when you buy any of these products
- medicines that can affect the acidity of your urine, such as vitamin C (ascorbic acid) or sodium bicarbonate (for example in medicines for indigestion)

If you are unsure about whether the medicines you are taking are in the list above, ask your doctor or pharmacist before taking [Product name].

Drug testing

This medicine may give a positive result when testing for drug use.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant, or are planning to have a baby, ask your doctor for advice before taking this medicine. Your body breaks down [Product name] into other substances which can cross the placenta and enter breast milk. Available data from the use of lisdexamfetamine dimesylate during the first three months of pregnancy do not indicate increased risk of congenital malformation in the child, but may increase the risk for pre-eclampsia (a condition usually occurring after 20 weeks of pregnancy characterized by high blood pressure and protein in the urine) and preterm birth. Newborns exposed to amphetamine during pregnancy may experience withdrawal symptoms (trembling, irritability, tight muscle tone).

You should not use this medicine during pregnancy unless explicitly advised by your doctor, or breast feed while taking [Product name].

Driving and using machines

You may feel dizzy, have problems focusing or have blurred vision when taking [Product name] If these happen it will be dangerous to do things such as drive or use machines.

[Product name] contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take [Product name]

How much to take

Always take this medicine exactly as described in this leaflet or as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Product name] is only for you. Do not give this medicine to anyone else, even if their symptoms seem similar.

The maximum daily dose is 70 mg.

How to take [Product name]

- Take [Product name] in the morning before breakfast. It can be taken with or without food.
- There are two ways to take [Product name]:
 - Swallow the capsule whole with a drink of water
 - Open the capsule and empty the contents into:

- Soft food such as yogurt
- A glass of water or orange juice

Use a spoon to break up any bits and stir the [Product name] and yogurt, water or orange juice until they are completely mixed together. Eat all the yogurt or drink all the water or orange juice immediately after mixing with [Product name]. Do not store it. Do not worry if there is a film left in the glass or container afterwards – this is not the active ingredient.

Dose

- Your doctor will tell you what strength of capsule to take ([Product name] 30 mg, 50 mg or 70 mg capsule) each day.
- The recommended dose at the start of treatment is 30 mg. Later on your doctor may increase this to 50 mg or 70 mg.
- If you have any kidney related problems your doctor may reduce the dose.

- Do not split the dose of a capsule; take the entire contents of the capsule. Do not take anything less than one capsule per day.

If you do not feel better after 1 month of treatment

If you do not feel better, tell your doctor. You may need a different treatment.

Not using [Product name] properly

- If [Product name] is not used properly, it may cause abnormal behaviour. You may also start to feel like you need the medicine for reasons other than helping with your ADHD. Tell your doctor if you have ever had a problem with alcohol, prescription medicines or street drugs

If you take more [Product name] than you should

If you take too much medicine, talk to a doctor or call an ambulance straight away. Tell them how much you have taken.

Signs of overdose may include: restlessness, shaking, increased uncontrolled movements, muscle twitching, fast breathing, being confused, an inclination to fight or quarrel, seeing, feeling or hearing things that are not real (hallucinations), panicked state, high fever or muscle breakdown. Tiredness and depression may follow. Being or feeling sick, vomiting, diarrhoea and stomach cramps may also occur. Changes in heartbeat (slow, fast or uneven), high or low blood pressure, circulatory collapse, fits and coma may be seen.

If you forget to take [Product name]

Do not take a double dose to make up for a forgotten dose. If you forget a dose, wait until the next day. Avoid taking it in the afternoon because of the possibility of sleep disturbances (insomnia).

If you stop taking [Product name]

If you stop taking this medicine, ADHD symptoms may come back.

Do not stop taking the medicine without first talking to your doctor. You should not suddenly stop taking this medicine on your own.

Things your doctor will do when you take [Product name]

Your doctor will do some tests

- before you start - to make sure that [Product name] is safe for you and will be of benefit to you.
- after you start - your doctor will do tests at least every 6 months, but possibly more often. The tests will also be done if the dose is changed. These tests will include:
 - checking your appetite
 - measuring weight
 - measuring blood pressure and heart rate
 - checking whether you have any problems with your mood, state of mind or any other unusual feelings, or if these have got worse while taking [Product name].

Long-term treatment

[Product name] does not need to be taken forever. If you take [Product name] for more than a year, your doctor should stop treatment for a short time. This will show if you still need the medicine.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Your doctor will talk to you about these side effects.

Some side effects could be serious. If you get any of the side effects below, see a doctor straight away:

Common (may affect up to 1 in 10 adults)

- uneven heartbeat (palpitations)
- chest pain (may be a sign of heart problems)

Uncommon (may affect up to 1 in 100 adults)

- feeling unusually excited, over-active, or un-inhibited (mania)
- allergic reaction (hypersensitivity)

Not known: frequency cannot be estimated from the available data

- seeing, feeling or hearing things that are not real (hallucinations)
- severe allergic reaction characterised by a sharp drop in blood pressure, difficulty breathing, and hives/itching (anaphylactic reaction)
- seeing or feeling or hearing things that are not real, paranoia, delusions(psychotic episodes)
- fits(seizures)
- abnormal heart rhythm, life-threatening irregular heart rhythm (seen on an electrocardiogram). See section 2, Warnings and precautions
- allergic liver injury seen as possible yellowing of the eyes and/or skin (eosinophilic hepatitis)
- swelling of the face, tongue or throat; difficulty swallowing; hives and breathing difficulties (angioedema) or serious skin rash , with or without blisters, of the skin and mucous membranes or skin irritation, sores or swelling in the mouth, throat, eyes, nose and around the genitals (Stevens Johnson syndrome)
- breathlessness or swelling of the legs (signs of heart muscle disease)

If you have any of the side effects above, see a doctor straight away.

Other side effects include the following. If they get serious, please tell your doctor or pharmacist:

Very common (may affect more than 1 in 10 adults)

- decreased appetite
- being unable to sleep
- dry mouth
- headache

Common (may affect up to 1 in 10 adults)

- feeling agitated, jittery, anxious, or irritable
- feeling unusually tired or restlessness
- unable to get or keep an erection or changes in sex drive
- feeling dizzy
- uncontrolled jerking, shaking, trembling or being unusually active
- hard-to-control, repeated twitching of any parts of the body or repeating sounds and words(tics)
- having mood swings
- high blood pressure, fast or uneven heartbeat (tachycardia)
- difficulty breathing
- feeling sick or diarrhoea
- constipation
- weight loss
- excessive sweating
- stomach pains
- grinding of the teeth

Uncommon (may affect up to 1 in 100 adults)

- talking excessively
- feeling depressed, anxious, low, or uneasy (dysphoria)
- feeling excessively happy or excited (euphoria)
- excessive picking of the skin
- uncontrolled twitching or jerking of the body
- feeling unusually sleepy
- itching, rash or raised red itchy rashes (urticaria)
- blurred vision
- high temperature (fever)
- being sick
- metallic taste or changes in taste (dysgeusia)
- fainting
- nosebleed

Not known: frequency cannot be estimated from the available data

- excessive widening of the pupils
- aggression
- poor blood circulation which makes the toes and fingers go numb and pale (Raynaud's phenomenon)

Effects on weight

[Product name] may cause weight loss in some people.

Your doctor will watch your weight, as well as how well you are eating. If you are losing weight, your doctor may stop treatment with [Product name] for a short time.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#).* By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [Product name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and the carton after EXP. The expiry date refers to the last day of that month.

Do not store above 30°C.

Do not use this medicine if the capsules look damaged in any way.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment

6. Contents of the pack and other information**What [Product name] contains**

The active substance is lisdexamfetamine dimesylate.

Each 30 mg capsule contains 30 mg lisdexamfetamine dimesylate, corresponding to 8.9 mg of dexamfetamine.

Each 50 mg capsule contains 50 mg lisdexamfetamine dimesylate, corresponding to 14.8 mg of dexamfetamine.

Each 70 mg capsule contains 70 mg lisdexamfetamine dimesylate, corresponding to 20.8 mg of dexamfetamine.

The other ingredients are:

- Capsule content: cellulose microcrystalline, croscarmellose sodium, magnesium stearate
- Capsule shell: gelatin, titanium dioxide (E171), black ink (shellac, propylene glycol, black iron oxide and potassium hydroxide)
- Capsules shell colourants:
 - 30 mg: yellow iron oxide (E172), red iron oxide (E172)
 - 50 mg: indigo carmine (E132).
 - 70 mg: yellow iron oxide (E172), red iron oxide (E172), indigo carmine (E132).

What [Product name] looks like and contents of the pack

Plastic bottle with child-resistant cap

30 mg: White body and orange cap, printed 'TEV' and '30 mg' in black ink.

50 mg: White body and blue cap, printed 'TEV' and '50 mg' in black ink.

70 mg: Blue body and orange cap, printed 'TEV' and '70 mg' in black ink.

Each capsule is approximately 19 mm long and 6 mm wide.

Pack sizes: 10 or 30 capsules.

Not all pack sizes may be marketed

Houder van de vergunning voor het in de handel brengen en fabrikant

Houder van de vergunning voor het in de handel brengen

Teva B.V.

Swensweg 5

2031 GA Haarlem

Nederland

Fabrikant

PLIVA Hrvatska d.o.o. (PLIVA Croatia LTD.)

Prilaz baruna Filipovića 25

10000 Zagreb

Kroatië

In het register ingeschreven onder

RVG 129375, harde capsules 30 mg

RVG 129376, harde capsules 50 mg

RVG 129377, harde capsules 70 mg

Dit medicijn is geregistreerd in lidstaten van de Europese Economische Ruimte onder de volgende namen:

Denemarken	Volidax
Finland	Volidax Adult 30 mg kapseli, kova Volidax Adult 50 mg kapseli, kova Volidax Adult 70 mg kapseli, kova
Duitsland	Lisdexamfetamin-ratiopharm Erwachsene 30 mg Hartkapseln Lisdexamfetamin-ratiopharm Erwachsene 50 mg Hartkapseln Lisdexamfetamin-ratiopharm Erwachsene 70 mg Hartkapseln
IJsland	Volidax
Nederland	Lisdexamfetamine Teva 30 mg, harde capsules Lisdexamfetamine Teva 50 mg, harde capsules Lisdexamfetamine Teva 70 mg, harde capsules
Noorwegen	Volidax
Zweden	Volidax

Deze bijsluiter is voor het laatst goedgekeurd in september 2023