

**PACKAGE LEAFLET**

## Package Leaflet: Information for the user

### Levonorgestrel Aristo 1.5 mg tabletten levonorgestrel

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Levonorgestrel Aristo is and what it is used for
2. What you need to know before you take Levonorgestrel Aristo
3. How to take Levonorgestrel Aristo
4. Possible side effects
5. How to store Levonorgestrel Aristo
6. Contents of the pack and other information

#### **1. What Levonorgestrel Aristo is and what it is used for**

Levonorgestrel Aristo is an emergency contraceptive that can be used within 72 hours (3 days) of unprotected sex or if your usual contraceptive method has failed.

It is about the following cases:

- No contraception was used during the sexual intercourse.
- The contraception measure was used incorrectly, for example as if a condom was penetrated, slipped away or used in a wrong way, if vaginal pessary or diaphragm changed position, burst, was broken or taken out ahead of time, in the case of a failed interruption during coitus interruptus (e.g. sperm ejaculated in vagina or on external genitalia).

This medicine contains a synthetic hormone-like active substance called levonorgestrel. It prevents about 84% of expected pregnancies when you take it within 72 hours of having unprotected sex. It will not prevent a pregnancy every time and is more effective if you take it as soon as possible after unprotected sex. It is better to take it within 12 hours rather than delay until the third day.

Levonorgestrel Aristo is thought to work by:

- stopping your ovaries from releasing an egg;
- preventing sperm from fertilising any egg you may have already released.

This medicine can only prevent you becoming pregnant if you take it within 72 hours of unprotected sex. It does not work if you are already pregnant. If another unprotected intercourse takes place after the use of Levonorgestrel Aristo (also if this is during the same menstrual cycle), the tablet will not exert its contraceptive effect and there is again the risk of pregnancy.

Levonorgestrel Aristo is not indicated for use before the first menstrual bleeding (menarche).

#### **2. What you need to know before you take Levonorgestrel Aristo**

##### **Do not take Levonorgestrel Aristo**

- If you are allergic to levonorgestrel or any of the other ingredients of this medicine listed in

section 6.

### **Warnings and precautions**

If any of the following applies to you, talk to your doctor before taking this medicine as emergency contraception may not be suitable for you. Your doctor may prescribe another type of emergency contraception for you.

- If you are pregnant or think that you may already be pregnant. This medicine will not work if you are already pregnant. If you are already pregnant, Levonorgestrel Aristo cannot terminate pregnancy, so this medicine is not an “abortion pill”.

You may already be pregnant if:

- your period is more than 5 days late, or you have experienced unusual bleeding when your next period is due
- you have had unprotected sex more than 72 hours ago, and since your last period

The use of this medicine is not advised if:

- you have a disease of your small bowel (such as Crohn’s disease) that inhibits the absorption of the medicine
- you have severe liver problems
- you have a history of ectopic pregnancy (where the baby develops somewhere outside the womb)
- you have ever had a disease called salpingitis (inflammation of the Fallopian tubes).

A previous ectopic pregnancy or previous infection of the fallopian tubes increase the risk of a new ectopic pregnancy.

In all women, emergency contraception should be taken as soon as possible after unprotected intercourse. There is some evidence that levonorgestrel (the active substance in this medicine) may be less effective with increasing body weight or body mass index (BMI), but these data were limited and inconclusive. Therefore, Levonorgestrel Aristo is still recommended for all women regardless of their weight or BMI.

If you are concerned about any problems related to taking emergency contraception talk with your doctor or other healthcare professional.

### **Children and adolescents**

This medicine is not indicated for use before the first menstrual bleeding (menarche).

### **If you are worried about sexually transmitted diseases**

If you did not use a condom (or if it has been torn or slid down) during the intercourse, it might be possible that you have caught a sexually transmitted disease or the HIV virus.

This medicine will not protect you against sexually transmitted diseases, only condoms can do this.

Ask your doctor, nurse, family planning clinic or pharmacist for advice if you are worried about this.

### **Other medicines and Levonorgestrel Aristo**

Tell your doctor or pharmacist if you are taking, have recently taken any other medicines, including medicines obtained without a prescription or herbal medicines.

Some medicines may prevent Levonorgestrel Aristo from working effectively. If you have used any of the medicines below during the last 4 weeks, this medicine may be less suitable for you. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD). If this is not an option for you or if you are unable to see your doctor promptly, you can take a double dose of Levonorgestrel Aristo.

- barbiturates and other medicines used to treat epilepsy (for example, primidone, phenytoin, and carbamazepine)
- medicines used to treat tuberculosis (for example, rifampicin, rifabutin)
- a treatment for HIV infection (ritonavir, efavirenz)
- a medicine used to treat fungal infections (griseofulvin)

- herbal remedies containing St John's wort (*Hypericum perforatum*).

Talk to your doctor or pharmacist if you need further advice on the correct dose for you.

Consult a doctor as soon as possible after taking the tablets for further advice on a reliable form of regular contraception and to exclude a pregnancy. (see also section 3, **How to take Levonorgestrel Aristo** for further advice).

Levonorgestrel Aristo may also affect how well other medicines work

- a medicine called cyclosporin (suppresses the immune system).

### **How often can you use Levonorgestrel Aristo**

You should only use Levonorgestrel Aristo in emergencies and not as a regular method of contraception. If this medicine is used more than once in a menstrual cycle, it is less reliable, and it is more likely to upset your menstrual cycle (period).

Levonorgestrel Aristo does not work as well as regular methods of contraception. Your doctor, practice nurse or family planning clinic can tell you about long-term methods of contraception which are more effective in preventing you from getting pregnant.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking any medicine.

#### ***Pregnancy***

You should not take this medicine if you are already pregnant. If you do become pregnant even after taking this medicine, it is important that you see your doctor. There is no evidence that levonorgestrel will harm a baby that develops in your uterus/ womb if you use this medicine as described.

Nevertheless, your doctor may want to check that the pregnancy is not ectopic (where the baby develops somewhere outside the womb). This is especially important if you develop severe abdominal pain after taking this medicine or if you have previously had an ectopic pregnancy, Fallopian tube surgery or pelvic inflammatory disease.

#### ***Breast-feeding***

The active ingredient of this medicine is excreted into your breast milk. Therefore, it is suggested that you take your tablet immediately after a breast-feeding and avoid nursing at least 8 hours following levonorgestrel administration then drain your milk with a breast pump for 8 hours following tablet taking. In this way you are taking your tablet well before the next feed and reducing the amount of active ingredient your baby may take in with the breast milk.

#### ***Fertility***

This medicine increases the possibility of menstruation disturbances which can sometimes lead to earlier or later ovulation date resulting in modified fertility date. Although there are no fertility data in the long term, after treatment with levonorgestrel a rapid return to fertility is expected and therefore, regular contraception should be continued or initiated as soon as possible after Levonorgestrel Aristo use.

### **Driving and using machines**

This medicine is unlikely to affect your ability to drive a car or use machines. However, if you feel tired or dizzy do not drive or operate machinery.

### **Levonorgestrel Aristo contains lactose (a type of sugar)**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### 3 How to take Levonorgestrel Aristo

Always take this medicine exactly as described in the leaflet or as your pharmacist or doctor has told you. Check with your doctor or pharmacist if you are not sure.

- Take the tablet as soon as possible, preferably within 12 hours, and no later than 72 hours (3 days) after you have had unprotected sex. Do not delay taking the tablet. The tablet works better the sooner you take it after having unprotected sex. It can only prevent you becoming pregnant, if you take it within 72 hours of unprotected sex.
- This medicine can be taken at any time in your menstrual cycle assuming you are not already pregnant or think you may be pregnant. Do not crush or chew the tablet. Swallow the tablet whole with water.
- If you are using one of the medicines that may prevent [Product Name[]] from working properly (see section above **Other medicines and Levonorgestrel Aristo**) or if you have used one of these medicines in the past 4 weeks, this medicine may work less effectively for you. Your doctor may prescribe another type of (non-hormonal) emergency contraceptive, i.e. a copper intrauterine device (Cu-IUD). If this is not an option for you or if you are unable to see your doctor promptly, you can take a double dose of Levonorgestrel Aristo (meaning 2 tablets taken together at the same time).
- If you are already using a regular method of contraception such as the contraceptive pill, you can continue to take this at your regular times.

If another unprotected intercourse takes place after the use of Levonorgestrel Aristo (also if this is during the same menstrual cycle), the tablet will not exert its contraceptive effect and there is again the risk of pregnancy.

If you are concerned about any problems related to taking emergency contraception talk with your doctor or other healthcare professional.

#### **What to do if you are sick (vomit)**

If you are sick (vomit) within three hours of taking the tablet, you should immediately take another tablet.

#### **After you have taken Levonorgestrel Aristo**

After you have taken this medicine, if you want to have sex and are not using the contraceptive pill, you should use condoms or a cap plus spermicide until your next menstrual period. This is because Levonorgestrel Aristo will not work if you have unprotected sex again, before your next period is due.

After you have taken this medicine, you are advised to make an appointment to see your doctor about three weeks later to make sure that the medicine has worked. If your period is more than 5 days late or is unusually light or unusually heavy, you should contact your doctor as soon as possible. If you do become pregnant even after taking this medicine, it is important that you see your doctor.

Your doctor can also tell you about longer-term methods of contraception which are more effective in preventing you from getting pregnant.

If you continue to use regular hormonal contraception such as the contraceptive pill and you do not have a bleed in your pill-free period, see your doctor to make sure you are not pregnant.

#### **Your next period after you took Levonorgestrel Aristo**

After the use of this medicine, your period is usually normal and will start at the usual day; however sometimes, this will be a few days later or earlier. If your period starts more than 5 days later than expected, an “abnormal” bleeding occurs at that time or if you think that you might be pregnant, you should check whether you are pregnant by a pregnancy test.

### **If you take more Levonorgestrel Aristo than you should**

Although there have been no reports of serious harmful effects from taking too many tablets at once, you may feel sick, actually be sick (vomit), or have vaginal bleeding. You should ask your pharmacist, doctor, practice nurse or family planning clinic for advice, especially if you have been sick, as the tablet may not have worked properly.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### **Very common (may affect more than 1 in 10 people):**

- Feeling sick (nausea)
- You might have some irregular bleeding until your next period
- You might have lower abdominal pain
- Tiredness
- Headache

### **Common (may affect up to 1 in 10 people):**

- Being sick (vomiting). If you are sick, read the section 'What to do if you are sick (vomit)'.  
- Your period might be different. Most women will have a normal period at the expected time, but some may have their period later or earlier than normal. You might also have some irregular bleeding or spotting until your next period. If your period is more than 5 days late or is unusually light or unusually heavy, you should contact your doctor as soon as possible.
- You might have tender breasts, diarrhoea, feel dizzy after taking this medicine.

### **Very rare effects (may affect up to 1 in 10,000 people):**

- Rash, urticaria, pruritus, swelling of the face, pelvic pain, painful period, abdominal pain.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

## **5. How to store Levonorgestrel Aristo**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton. The expiry date refers to the last day of the month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Levonorgestrel Aristo contains**

The active substance is levonorgestrel. Each tablet contains 1.5 milligrams of levonorgestrel.

The other ingredients are: potato starch, maize starch, silica, colloidal anhydrous (E 551), magnesium stearate (E 470b), talc (E 553b) and lactose monohydrate.

### **What Levonorgestrel Aristo looks like and contents of the pack**

Levonorgestrel Aristo 1.5 mg tabletten is a white to off white, round bevel edged, flat faced tablet debossed with “J06” on one side and plain on other side.

Levonorgestrel Aristo is available in clear and transparent PVC/Aluminium-blister containing one tablet. The blister is packaged in a folded carton.

### **Marketing Authorisation Holder**

Aristo Pharma GmbH Ltd.  
Berlin  
Wallenroder Strasse  
8-10, 13435 Berlin

### **Manufacturer**

Pharmadox Healthcare Ltd.  
KW20A Kordin Industrial Park,  
Paola PLA 3000, Malta.

Aristo Pharma GmbH  
Wallenroder Strasse 8-10,  
13435 Berlin  
Germany

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Nederland:	Levonorgestrel Aristo 1,5 mg tabletten
Duitsland:	Levonorgestrel Adalvo 1.5 mg Tabletten
Italië:	Levonorgestrel Adalvo
Spanje:	Levonorgestrel Adalvo 1.5 mg comprimidos EFG

[To be completed nationally]

**This leaflet was last revised in April 2026.**