

Package leaflet: information for the user

Macitentan CF 10 mg, filmomhulde tabletten macitentan

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What <Product name> is and what it is used for
2. What you need to know before you take <Product name>
3. How to take <Product name>
4. Possible side effects
5. How to store <Product name>
6. Contents of the pack and other information

1. What <Product name> is and what it is used for

<Product name> contains the active substance macitentan, which belongs to the class of medicines called “endothelin receptor antagonists”.

<Product name> is used for the long-term treatment of pulmonary arterial hypertension (PAH):

- in adults of WHO Functional Class (FC) II to III
- in children under 18 years and body weight of at least 40 kg with WHO Functional Class (FC) II to III

It can be used on its own or with other medicines for PAH. PAH is high blood pressure in the blood vessels that carry blood from the heart to the lungs (the pulmonary arteries). In people with PAH, these arteries get narrower, so the heart has to work harder to pump blood through them. This causes people to feel tired, dizzy, and short of breath.

<Product name> widens the pulmonary arteries, making it easier for the heart to pump blood through them. This lowers the blood pressure, relieves the symptoms and improves the course of the disease.

2. What you need to know before you take <Product name>

DO NOT take <Product name>

- if you are allergic to macitentan or any of the other ingredients of this medicine (listed in section 6)
- if you are pregnant, if you are planning to become pregnant, or if you could become pregnant because you are not using reliable birth control (contraception). See section ‘Pregnancy and breast-feeding’
- if you are breast-feeding. See section ‘Pregnancy and breast-feeding’
- if you have liver disease or if you have very high levels of liver enzymes in your blood. Talk to your doctor, who will decide whether this medicine is suitable for you

If any of these apply to you, please tell your doctor.

Warnings and precautions

Talk to your doctor or pharmacist before taking <Product name>.

You will need blood tests, as indicated by your doctor:

Your doctor will take blood test before you start treatment with <Product name> and during treatment to test:

- whether you have anaemia (a reduced number of red blood cells)
- whether your liver is working properly

If you have anaemia (a reduced number of red blood cells), you may have the following signs:

- dizziness
- fatigue/malaise/weakness
- fast heart rate, palpitations
- pallor

If you notice any of these signs, **tell your doctor**.

Signs that your liver may not be working properly include:

- feeling sick (nausea)
- vomiting
- fever
- pain in your stomach (abdomen)
- yellowing of your skin or the whites of your eyes (jaundice)
- dark-coloured urine
- itching of your skin
- unusual tiredness or exhaustion (lethargy or fatigue)
- flu-like syndrome (joint and muscle pain with fever)

If you notice any of these signs, **tell your doctor immediately**.

If you have kidney problems, talk to your doctor before using <Product name>. Macitentan may lead to more reduction of blood pressure and decrease in haemoglobin in patients with kidney problems.

In patients with pulmonary veno-occlusive disease (obstruction of the lung veins), the use of medicines for treatment of PAH, including <Product name>, may lead to pulmonary oedema. If you have signs of pulmonary oedema when using <Product name>, such as a sudden, important increase in breathlessness and low oxygen, **tell your doctor immediately**. Your doctor may perform additional tests, and will determine what treatment regimen is most suitable for you.

Children and adolescents

Do not give this medicine to children below 2 years of age because efficacy and safety have not been established.

Other medicines and <Product name>

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicine. <Product name> can affect other medicines.

If you take <Product name> together with other medicines including those listed below, the effects of <Product name> or the other medicines might be altered. Please talk to your doctor or pharmacist if you are taking any of the following medicines:

- rifampicin, clarithromycin, telithromycin, ciprofloxacin, erythromycin (antibiotics used to treat infections)

- phenytoin (a medicine used to treat seizures)
- carbamazepine (used to treat depression and epilepsy)
- St. John's Wort (an herbal preparation used to treat depression)
- ritonavir, saquinavir (used to treat HIV infections)
- nefazodone (used to treat depression)
- ketoconazole (except shampoo), fluconazole, itraconazole, miconazole, voriconazole (medicines used against fungal infections)
- amiodarone (to control the heartbeat)
- cyclosporine (used to prevent organ rejection after transplant)
- diltiazem, verapamil (to treat high blood pressure or specific heart problems)

<Product name> with food

If you are taking piperine as a dietary supplement, this may alter how the body responds to some medicinal products, including <Product name>. Please talk to your doctor or pharmacist should this be the case.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

<Product name> may harm unborn babies conceived before, during or soon after treatment.

- If it is possible you could become pregnant, use a reliable form of birth control (contraception) while you are taking <Product name>. Talk to your doctor about this.
- Do not take <Product name> if you are pregnant or planning to become pregnant.
- If you become pregnant or think that you may be pregnant while you are taking <Product name>, or shortly after stopping <Product name> (up to 1 month), see your doctor immediately.

If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you start taking <Product name> and regularly (once a month) while you are taking <Product name>.

It is not known if macitentan is transferred to breast milk. Do not breast-feed while you are taking <Product name>. Talk to your doctor about this.

Fertility

If you are a man taking <Product name>, it is possible that this medicine may lower your sperm count. Talk to your doctor if you have any questions or concerns about this.

Driving and using machines

<Product name> can cause side effects such as headaches and hypotension (listed in section 4), and the symptoms of your condition can also make you less fit to drive or use machines.

<Product name> contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take <Product name>

<Product name> should only be prescribed by a doctor experienced in the treatment of pulmonary arterial hypertension.

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Adults and children aged less than 18 years weighing at least 40 kg

The recommended dose of <Product name> is one 10 mg tablet, once a day. Swallow the whole tablet, with a glass of water. Do not chew or break the tablet in order to avoid damage to the film-coating. You can take <Product name> with or without food. It is best to take the tablet at the same time each day.

For children weighing less than 40 kg, macitentan dispersible tablets with a lower strength are available under other brand names.

If you take more <Product name> than you should

If you have taken more tablets than you have been told to take, you may experience headache, nausea, or vomiting. Ask your doctor for advice.

If you forget to take <Product name>

If you forget to take <Product name>, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for a forgotten tablet.

If you stop taking <Product name>

<Product name> is a treatment that you will need to keep on taking to control your PAH. Do not stop taking <Product name> unless you have agreed this with your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Uncommon serious side effects (may affect up to 1 in 100 people)

- Allergic reactions (swelling around the eyes, face, lips, tongue or throat, itching and/or rash)
- If you notice any of these signs, tell your doctor immediately.

Very common side effects (may affect more than 1 in 10 people)

- Anaemia (low number of red blood cells) or reduced haemoglobin
- Headache
- Bronchitis (inflammation of the airways)
- Nasopharyngitis (inflammation of the throat and nasal passages)
- Oedema (swelling), especially of the ankles and feet

Common side effects (may affect up to 1 in 10 people)

- Pharyngitis (inflammation of the throat)
- Influenza (flu)
- Urinary tract infection (bladder infection)
- Hypotension (low blood pressure)
- Nasal congestion (blocked nose)
- Elevated liver tests
- Leukopenia (decreased white blood cell counts)
- Thrombocytopenia (decreased blood platelet counts)
- Flushing (redness of the skin)
- Increased uterine bleeding

Side effects in children and adolescents

The side effects listed above may also be seen in children. Additional side effects very commonly seen in children include upper respiratory tract infection (infected nose sinuses, or

throat) and gastroenteritis (inflamed stomach and gut). Rhinitis (itchy, runny, or blocked nose) was seen commonly in children.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store <Product name>

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer require. These measures will help protect the environment.

6. Contents of the pack and other information

What <Product name> contains

The active substance is macitentan. Each film-coated tablet contains 10 mg macitentan.

The other ingredients are maltodextrin (E1400), polysorbate 80 (E433), mannitol (E421), starch, pregelatinised (maize starch), croscarmellose sodium (E468), magnesium stearate (E572), poly(vinyl alcohol) (E1203), talc (E553b), titanium dioxide (E171), glycerol monocaprylocaprate, type I, sodium laurilsulfate (E487).

What <Product name> looks like and contents of the pack

<Product Name> is a film-coated tablet.

White to off-white, round, biconvex film-coated tablets with a diameter of approximately 5.6 mm.

<Product name> 10 mg is available in PVC/PE/PVdC-Al blisters containing 15 or 30 film-coated tablets or PVC/PE/PVdC-Al unit-dose blisters containing 15x1 or 30x1 film-coated tablets.

Not all pack sizes may be marketed.

Houder van de vergunning voor het in de handel brengen en fabrikant

Vergunninghouder:

Centrafarm B.V.

Van de Reijtstraat 31-E

4814 NE Breda

Nederland

Fabrikant:

STADA Arzneimittel AG

Stadastrasse 2 - 18
61118 Bad Vilbel
Duitsland

STADA Arzneimittel GmbH
Muthgasse 36/2
1190 Wenen
Oostenrijk

Clonmel Healthcare Ltd
Waterford Road, Clonmel,
E91 D768 Co. Tipperary
Ierland

Centrafarm Services B.V.
Van de Reijtstraat 31 E
4814NE Breda
Nederland

In het register ingeschreven onder

Macitentan CF 10 mg, filmomhulde tabletten RVG 132854

Dit medicijn is geregistreerd in lidstaten van de Europese Economische Ruimte onder de volgende namen:

België, Italië, Luxemburg:	Macitentan EG
Denemarken, Duitsland, Finland, Hongarije, IJsland, Noorwegen, Oostenrijk, Polen, Portugal, Roemenië, Slowakije, Spanje, Zweden:	Macitentan STADA
Ierland, Malta:	Macitentan Clonmel
Nederland:	Macitentan CF

Deze bijsluiter is voor het laatst goedgekeurd in juli 2025.