

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAAM VAN HET GENEESMIDDEL

Ezetimibe/Atorvastatine GL 10 mg/ 10 mg, tabletten
Ezetimibe/Atorvastatine GL 10 mg/ 20 mg, tabletten
Ezetimibe/Atorvastatine GL 10 mg/ 40 mg, tabletten
Ezetimibe/Atorvastatine GL 10 mg/ 80 mg, tabletten

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 10 mg of ezetimibe and 10 mg, 20 mg, 40 mg or 80 mg of atorvastatin (as atorvastatin calcium trihydrate).

Excipients with known effect

Each 10 mg/10 mg tablet contains 145 mg lactose.
Each 10 mg/20 mg tablet contains 170 mg lactose.
Each 10 mg/40 mg tablet contains 219 mg lactose.
Each 10 mg/80 mg tablet contains 317 mg lactose.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablet.

10 mg/10 mg tablet: White to off white, capsule shaped tablets (12.7 mm x 5.1 mm) on one side debossed with "1"

10 mg/20 mg tablet: White to off white, capsule shaped tablets (14.5 mm x 6.8 mm) on one side debossed with "2"

10 mg/40 mg tablet: White to off white, capsule shaped tablets (16.4 mm x 6.3 mm) on one side debossed with "3"

10 mg/80 mg tablet: White to off white, capsule shaped tablets (17.0 mm x 8.0 mm) on one side debossed with "4"

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Ezetimibe/Atorvastatine GL as an adjunct to diet is indicated as substitution therapy for treatment of adults with primary (heterozygous and homozygous familial and non-familial) hypercholesterolaemia or mixed hyperlipidaemia already controlled with atorvastatin and ezetimibe given concurrently at the same dose level.

4.2 Posology and method of administration

Posology

The recommended dose of Ezetimibe/Atorvastatine GL is 1 tablet per day.

The maximum recommended dose of Ezetimibe/Atorvastatine GL is 10 mg/80 mg per day. The patient should be on an appropriate lipid lowering diet and should continue on this diet during treatment with Ezetimibe/Atorvastatine GL.

Ezetimibe/Atorvastatine GL is not suitable for initial therapy. Treatment initiation or dose adjustment if necessary should only be done with the monocomponents and after setting the appropriate doses the switch to the fixed dose combination of the appropriate strength is possible.

Elderly

No dose adjustment is required for elderly patients (see section 5.2).

Paediatric population

The safety and efficacy of Ezetimibe/Atorvastatine GL in children and adolescent has not been established (see section 5.2). No data are available.

Hepatic impairment

Ezetimibe/Atorvastatine GL should be used with caution in patients with hepatic impairment (see sections 4.4. and 5.2). Ezetimibe/Atorvastatine GL is contraindicated in patients with active liver disease (see section 4.3).

Renal impairment

No dose adjustment is required for renally impaired patients (see section 5.2).

Coadministration with bile acid sequestrants

Dosing of Ezetimibe/Atorvastatine GL should occur either ≥ 2 hours before or ≥ 4 hours after administration of a bile acid sequestrant.

Coadministration with other medicines

In patients taking hepatitis C antiviral agents elbasvir/grazoprevir concomitantly with atorvastatin, the dose of atorvastatin should not exceed 20 mg/day (see section 4.4. and 4.5.).

Method of administration

Ezetimibe/Atorvastatine GL is for oral administration. The tablet should be swallowed with a sufficient amount of fluid (e. g. one glass of water).

Ezetimibe/Atorvastatine GL can be administered as a single dose at any (but preferable always the same) time of the day, with or without food.

4.3 Contraindications

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

Therapy with Ezetimibe/Atorvastatine GL is contraindicated during pregnancy and breast-feeding, and in women of child-bearing potential not using appropriate contraceptive measures (see section 4.6).

Ezetimibe/Atorvastatine GL is contraindicated in patients with active liver disease or unexplained persistent elevations in serum transaminases exceeding 3 times the upper limit of normal (ULN) and in patients treated with the hepatitis C antivirals glecaprevir/pibrentasvir.

4.4 Special warnings and precautions for use

Myopathy/Rhabdomyolysis

In post-marketing experience with ezetimibe, cases of myopathy and rhabdomyolysis have been reported. Most patients who developed rhabdomyolysis were taking a statin concomitantly with ezetimibe. However, rhabdomyolysis has been reported very rarely with ezetimibe monotherapy and

very rarely with the addition of ezetimibe to other agents known to be associated with increased risk of rhabdomyolysis.

Ezetimibe/Atorvastatine GL contains atorvastatin. Atorvastatin, like other HMG-CoA reductase inhibitors, may in rare occasions affect the skeletal muscle and cause myalgia, myositis, and myopathy that may progress to rhabdomyolysis, a potentially life-threatening condition characterised by markedly elevated creatine phosphokinase (CPK) levels (>10 times ULN), myoglobinaemia and myoglobinuria, which may lead to renal failure.

There have been very rare reports of an immune-mediated necrotising myopathy (IMNM) during or after treatment with statins, including atorvastatin. IMNM is clinically characterised by persistent proximal muscle weakness and elevated serum creatine kinase, which persist despite discontinuation of statin treatment.

Before the treatment

Ezetimibe/Atorvastatine GL should be prescribed with caution to patients with pre-disposing factors for rhabdomyolysis. ACPK level should be measured before starting treatment in the following situations:

- renal impairment
- hypothyroidism
- personal or familial history of hereditary muscular disorders
- previous history of muscular toxicity with a statin or fibrate
- previous history of liver disease and/or where substantial quantities of alcohol are consumed
- in elderly (age > 70 years), the necessity of such measurement should be considered, according to the presence of other predisposing factors for rhabdomyolysis
- situations where an increase in plasma levels may occur, such as interactions (see section 4.5) and special populations including genetic subpopulations (see section 5.2)

In such situations, the risk of treatment should be considered in relation to possible benefit, and clinical monitoring is recommended.

If CPK levels are significantly elevated (> 5 times ULN) at baseline, treatment should not be started.

Creatine phosphokinase measurement

Creatine phosphokinase (CPK) should not be measured following strenuous exercise or in the presence of any other cause of CPK increase as it makes value interpretation difficult. If CPK levels are significantly elevated at baseline (> 5 times ULN), levels should be remeasured within 5 to 7 days later to confirm the results.

Monitoring during treatment

- Patients must be asked to promptly report muscle pain, cramps, or weakness especially if accompanied by malaise or fever or if muscle signs and symptoms persist after discontinuing Ezetimibe/Atorvastatine GL.
- If such symptoms occur whilst a patient is receiving treatment with Ezetimibe/Atorvastatine GL, their CPK levels should be measured. If these levels are found to be significantly elevated (> 5 times ULN), treatment should be stopped.
- If muscular symptoms are severe and cause daily discomfort, even if the CPK levels are elevated to ≤ 5 times ULN, treatment discontinuation should be considered.
- If symptoms resolve and CPK levels return to normal, then re-introduction of Ezetimibe/Atorvastatine GL or introduction of another statin-containing product may be considered at the lowest dose and with close monitoring.
- Ezetimibe/Atorvastatine GL must be discontinued if clinically significant elevation of CPK levels (> 10 times ULN) occur, or if rhabdomyolysis is diagnosed or suspected.

Simultaneous treatment with other products

Due to the atorvastatin component of Ezetimibe/Atorvastatine GL, the risk of rhabdomyolysis is increased when atorvastatin is administered concomitantly with certain medicinal products that may increase the plasma concentration of atorvastatin such as potent inhibitors of CYP3A4 or transport proteins (e.g. ciclosporin, telithromycin, clarithromycin, delavirdine, stiripentol, ketoconazole, voriconazole, itraconazole, posaconazole, and certain HIV protease inhibitors including ritonavir, lopinavir, atazanavir, indinavir, darunavir, tipranavir/ritonavir etc). The risk of myopathy may also be increased with the concomitant use of gemfibrozil and other fibric acid derivatives, antivirals for the treatment of hepatitis C (HCV) (boceprevir, telaprevir, elbasvir/grazoprevir), erythromycin, or niacin. If possible, alternative (non-interacting) therapies should be considered instead of these medicinal products (see section 4.8).

In cases where coadministration of these medicinal products with Ezetimibe/Atorvastatine GL is necessary, the benefit and the risk of concurrent treatment should be carefully considered. When patients are receiving medicinal products that increase the plasma concentration of atorvastatin, a lower maximum dose of Ezetimibe/Atorvastatine GL is recommended. In addition, in the case of potent CYP3A4 inhibitors, a lower starting dose of Ezetimibe/Atorvastatine GL should be considered and appropriate clinical monitoring of these patients is recommended (see section 4.5).

Ezetimibe/Atorvastatine GL must not be co-administered with systemic formulations of fusidic acid or within 7 days of stopping fusidic acid treatment. In patients where the use of systemic fusidic acid is considered essential, statin treatment should be discontinued throughout the duration of fusidic acid treatment. There have been reports of rhabdomyolysis (including some fatalities) in patients receiving fusidic acid and statins in combination (see section 4.5). The patient should be advised to seek medical advice immediately if they experience any symptoms of muscle weakness, pain or tenderness

Statin therapy may be re-introduced seven days after the last dose of fusidic acid.

In exceptional circumstances, where prolonged systemic fusidic acid is needed, e.g. for the treatment of severe infection, the need for co-administration of Ezetimibe/Atorvastatine GL and fusidic acid should only be considered on a case by case basis and under close medical supervision.

Daptomycin

Cases of myopathy and/or rhabdomyolysis have been reported with HMG-CoA reductase inhibitors (e.g. atorvastatin and ezetimibe/atorvastatin) co-administered with daptomycin. Caution should be used when prescribing HMG-CoA reductase inhibitors with daptomycin, as either agent can cause myopathy and/or rhabdomyolysis when given alone. Consideration should be given to temporarily suspend Ezetimibe/Atorvastatine GL in patients taking daptomycin unless the benefits of concomitant administration outweigh the risk. Consult the prescribing information of daptomycin to obtain further information about this potential interaction with HMG-CoA reductase inhibitors (e.g. atorvastatin and ezetimibe/atorvastatin) and for further guidance related to monitoring (see section 4.5).

Liver Enzymes

In controlled coadministration trials in patients receiving ezetimibe and a statin, consecutive transaminase elevations (≥ 3 times the upper limit of normal [ULN]) have been observed (see section 4.8).

Liver function tests should be performed before the initiation of treatment and periodically thereafter. Patients who develop any signs or symptoms suggestive of liver injury should have liver function tests performed. Patients who develop increased transaminase levels should be monitored until the abnormality(ies) resolve. Should an increase in transaminase of greater than 3 times the ULN persist, reduction of dose or withdrawal of Ezetimibe/Atorvastatine GL is recommended.

Ezetimibe/Atorvastatine GL should be used with caution in patients who consume substantial quantities of alcohol and/or have a history of liver disease.

Hepatic Insufficiency

Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate or severe hepatic insufficiency, the administration of Ezetimibe/Atorvastatine GL is not recommended (see section 5.2).

Fibrates

The safety and efficacy of ezetimibe administered with fibrates have not been established. Therefore, concomitant treatment with Ezetimibe/Atorvastatine GL is not recommended (see section 4.5).

Ciclosporin

Caution should be exercised when initiating Ezetimibe/Atorvastatine GL in the setting of ciclosporin. Ciclosporin concentrations should be monitored in patients receiving Ezetimibe/Atorvastatine GL and ciclosporin (see section 4.5).

Anticoagulants

If Ezetimibe/Atorvastatine GL is added to warfarin, another coumarin anticoagulant, or fludione, the International Normalised Ratio (INR) should be appropriately monitored (see section 4.5).

Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL)

In a post-hoc analysis of stroke subtypes in patients without coronary heart disease (CHD) who had a recent stroke or transient ischemic attack (TIA) there was a higher incidence of hemorrhagic stroke in patients initiated on atorvastatin 80 mg compared to placebo. The increased risk was particularly noted in patients with prior hemorrhagic stroke or lacunar infarct at study entry. For patients with prior hemorrhagic stroke or lacunar infarct, the balance of risks and benefits of atorvastatin 80 mg is uncertain, and the potential risk of hemorrhagic stroke should be carefully considered before initiating treatment (see section 5.1).

Interstitial lung disease

Exceptional cases of interstitial lung disease have been reported with some statins, especially with long term therapy (see section 4.8). Presenting features can include dyspnoea, non-productive cough and deterioration in general health (fatigue, weight loss and fever). If it is suspected a patient has developed interstitial lung disease, statin therapy should be discontinued.

Diabetes mellitus

Some evidence suggests that statins as a class raise blood glucose and in some patients, at high risk of future diabetes, may produce a level of hyperglycaemia where formal diabetes care is appropriate. This risk, however, is outweighed by the reduction in vascular risk with statins and therefore should not be a reason for stopping statin treatment. Patients at risk (fasting glucose 5.6 to 6.9 mmol/L, BMI > 30 kg/m², raised triglycerides, hypertension) should be monitored both clinically and biochemically according to national guidelines.

Myasthenia

In few cases, statins have been reported to induce *de novo* or aggravate pre-existing myasthenia gravis or ocular myasthenia (see section 4.8). Ezetimibe/Atorvastatine GL should be discontinued in case of aggravation of symptoms. Recurrences when the same or a different statin was (re-)administered have been reported.

Excipients

Ezetimibe/Atorvastatine GL contains lactose.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency, or glucose-galactose malabsorption should not take this medicine.

Ezetimibe/Atorvastatine GL contains sodium.

Ezetimibe/Atorvastatine GL contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium free'.

4.5 Interaction with other medicinal products and other forms of interactionPharmacodynamic interactions

Multiple mechanisms may contribute to potential interactions with HMG Co-A reductase inhibitors. Drugs or herbal products that inhibit certain enzymes (e.g CYP3A4) and/or transporter (e.g., OATP1B) pathways may increase atorvastatin plasma concentrations and may lead to an increased risk of myopathy/rhabdomyolysis.

Consult the prescribing information of all concomitantly used drugs to obtain further information about their potential interactions with atorvastatin and/or the potential for enzyme or transporter alterations and possible adjustments to dose and regimens.

Pharmacokinetic interactions

No clinically significant pharmacokinetic interaction was seen when ezetimibe was coadministered with atorvastatin.

Effects of other medicinal products on Ezetimibe/Atorvastatine GL*Ezetimibe*

Antacids: Concomitant antacid administration decreased the rate of absorption of ezetimibe but had no effect on the bioavailability of ezetimibe. This decreased rate of absorption is not considered clinically significant.

Cholestyramine: Concomitant cholestyramine administration decreased the mean area under the curve (AUC) of total ezetimibe (ezetimibe + ezetimibe glucuronide) approximately 55%. The incremental low-density lipoprotein cholesterol (LDL-C) reduction due to adding ezetimibe to cholestyramine may be lessened by this interaction (see section 4.2).

Ciclosporin: In a study of eight post-renal transplant patients with creatinine clearance of > 50 mL/min on a stable dose of ciclosporin, a single 10 mg dose of ezetimibe resulted in a 3.4-fold (range 2.3- to 7.9-fold) increase in the mean AUC for total ezetimibe compared to a healthy control population, receiving ezetimibe alone, from another study (n=17). In a different study, a renal transplant patient with severe renal insufficiency who was receiving ciclosporin and multiple other medicinal products demonstrated a 12-fold greater exposure to total ezetimibe compared to concurrent controls receiving ezetimibe alone. In a two-period crossover study in twelve healthy subjects, daily administration of 20 mg ezetimibe for 8 days with a single 100-mg dose of ciclosporin on Day 7 resulted in a mean 15% increase in ciclosporin AUC (range 10% decrease to 51% increase) compared to a single 100-mg dose of ciclosporin alone. A controlled study on the effect of coadministered ezetimibe on ciclosporin exposure in renal transplant patients has not been conducted. Caution should be exercised when initiating Ezetimibe/Atorvastatine GL in the setting of ciclosporin. Ciclosporin concentrations should be monitored in patients receiving Ezetimibe/Atorvastatine GL and ciclosporin (see section 4.4).

Fibrates: Concomitant fenofibrate or gemfibrozil administration modestly increased total ezetimibe concentrations approximately 1.5- and 1.7-fold, respectively. Although these increases are not considered clinically significant, coadministration of Ezetimibe/Atorvastatine GL with fibrates is not recommended (see section 4.4).

Atorvastatin

Effect of co-administered medicinal products on atorvastatin

Atorvastatin is metabolized by cytochrome P450 3A4 (CYP3A4) and is a substrate of the hepatic transporters, organic anion-transporting polypeptide 1B1 (OATP1B1) and 1B3 (OATP1B3) transporter. Metabolites of atorvastatin are substrates of OATP1B1. Atorvastatin is also identified as a substrate of the multi-drug resistance protein 1 (MDR1) and breast cancer resistance protein (BCRP), which may limit the intestinal absorption and biliary clearance of atorvastatin (see section 5.2). Concomitant administration of medicinal products that are inhibitors of CYP3A4 or transport proteins may lead to increased plasma concentrations of atorvastatin and an increased risk of myopathy. The risk might also be increased at concomitant administration of atorvastatin with other medicinal products that have a potential to induce myopathy, such as fibric acid derivatives and ezetimibe (see section 4.3 and 4.4).

CYP3A4 inhibitors: Potent CYP3A4 inhibitors have been shown to lead to markedly increased concentrations of atorvastatin (see Table 1 and specific information below). Coadministration of potent CYP3A4 inhibitors (e.g. ciclosporin, telithromycin, clarithromycin, delavirdine, stiripentol, ketoconazole, voriconazole, itraconazole, posaconazole, some antivirals used in the treatment of HCV (e.g. elbasvir/grazoprevir) and HIV protease inhibitors including ritonavir, lopinavir, atazanavir, indinavir, darunavir, etc.) should be avoided if possible. In cases where coadministration of these medicinal products with atorvastatin cannot be avoided, lower starting and maximum doses of atorvastatin should be considered and appropriate clinical monitoring of the patient is recommended (see Table 1).

Moderate CYP3A4 inhibitors (e.g. erythromycin, diltiazem, verapamil and fluconazole) may increase plasma concentrations of atorvastatin (see Table 1). An increased risk of myopathy has been observed with the use of erythromycin in combination with statins. Interaction studies evaluating the effects of amiodarone or verapamil on atorvastatin have not been conducted. Both amiodarone and verapamil are known to inhibit CYP3A4 activity and coadministration with Ezetimibe/Atorvastatine GL may result in increased exposure to atorvastatin. Therefore, a lower maximum dose of Ezetimibe/Atorvastatine GL should be considered and appropriate clinical monitoring of the patient is recommended when concomitantly used with moderate CYP3A4 inhibitors. Appropriate clinical monitoring is recommended after initiation or following dose adjustments of the inhibitor.

Inhibitors of Breast Cancer Resistant Protein (BCRP): Concomitant administration of products that are inhibitors of BCRP (e.g., elbasvir and grazoprevir) may lead to increased plasma concentration of atorvastatin and an increased risk of myopathy; therefore, a dose adjustment of atorvastatin should be considered depending on the prescribed dose. Co-administration of elbasvir and grazoprevir with atorvastatin increases plasma concentrations of atorvastatin 1.9-fold (see Table 1); therefore, the dose of Ezetimibe/Atorvastatine GL should not exceed 10 mg/20 mg daily in patients receiving concomitant medications with products containing elbasvir or grazoprevir (see sections 4.2 and 4.4).

Inducers of cytochrome P450 3A4: Concomitant administration of atorvastatin with inducers of cytochrome P450 3A4 (e.g., efavirenz, rifampicin, St. John's Wort) can lead to variable reductions in plasma concentrations of atorvastatin. Due to the dual interaction mechanism of rifampicin, (cytochrome P450 3A4 induction and inhibition of hepatocyte uptake transporter OATP1B1), simultaneous coadministration of atorvastatin with rifampicin is recommended, as delayed administration of atorvastatin after administration of rifampicin has been associated with a significant reduction in atorvastatin plasma concentrations. The effect of rifampicin on atorvastatin concentrations in hepatocytes is, however, unknown and if concomitant administration cannot be avoided, patients should be carefully monitored for efficacy.

Transport inhibitors: Inhibitors of transport proteins (e.g. ciclosporin) can increase the systemic exposure of atorvastatin (see Table 1). The effect of inhibition of hepatic uptake transporters on atorvastatin concentrations in hepatocytes is unknown. If concomitant administration cannot be avoided, a dose reduction of Ezetimibe/Atorvastatine GL and clinical monitoring for efficacy is recommended (see table 1).

Gemfibrozil/fibric acid derivatives: The use of fibrates alone is occasionally associated with muscle-related events, including rhabdomyolysis. The risk of these events may be increased with the concomitant use of fibric acid derivatives and atorvastatin.

Ezetimibe: The use of ezetimibe alone is associated with muscle-related events, including rhabdomyolysis. The risk of these events may therefore be increased with concomitant use of ezetimibe and atorvastatin. Appropriate clinical monitoring of these patients is recommended.

Colestipol: Plasma concentrations of atorvastatin and its active metabolites were lower when colestipol was coadministered with atorvastatin (by approx.. 25%). However, the lipid decreasing effects were greater when atorvastatin and colestipol were coadministered than when either medicinal product was given alone.

Fusidic acid: The risk of myopathy including rhabdomyolysis may be increased by the concomitant administration of systemic fusidic acid with statins. The mechanism of this interaction (whether it is pharmacodynamic or pharmacokinetic, or both) is yet unknown. There have been reports of rhabdomyolysis (including some fatalities) in patients receiving this combination. If treatment with systemic fusidic acid is necessary, atorvastatin treatment should be discontinued throughout the duration of the fusidic acid treatment (see section 4.4).

Colchicine: Although interaction studies with atorvastatin and colchicine have not been conducted, cases of myopathy have been reported with atorvastatin co-administered with colchicine, and caution should be exercised when prescribing atorvastatin with colchicine.

Daptomycin: The risk of myopathy and/or rhabdomyolysis may be increased by concomitant administration of HMG-CoA reductase inhibitors and daptomycin. Consideration should be given to suspending Ezetimibe/Atorvastatine GL temporarily in patients taking daptomycin unless the benefits of concomitant administration outweigh the risk (see section 4.4).

Boceprevir: Exposure to atorvastatin was increased when administered with boceprevir. When coadministration with Ezetimibe/Atorvastatine GL is required, starting with the lowest possible dose of Ezetimibe/Atorvastatine GL should be considered with titration up to desired clinical effect while monitoring for safety, without exceeding a daily dose of 10/20 mg. For patients currently taking Ezetimibe/Atorvastatine GL, the dose of Ezetimibe/Atorvastatine GL should not exceed a daily dose of 10 mg/20 mg during coadministration with boceprevir.

Effects of Ezetimibe/Atorvastatine GL on the pharmacokinetics of other medicinal products

Ezetimibe

In preclinical studies, it has been shown that ezetimibe does not induce cytochrome P450 drug metabolising enzymes. No clinically significant pharmacokinetic interactions have been observed between ezetimibe and drugs known to be metabolised by cytochromes P450 1A2, 2D6, 2C8, 2C9, and 3A4, or N-acetyltransferase.

Anticoagulants: Concomitant administration of ezetimibe (10 mg once daily) had no significant effect on bioavailability of warfarin and prothrombin time in a study of twelve healthy adult males. However, there have been post-marketing reports of increased International Normalised Ratio (INR) in patients who had ezetimibe added to warfarin or fluindione. If Ezetimibe/Atorvastatine GL is added to

warfarin, another coumarin anticoagulant, or fluindione, INR should be appropriately monitored (see section 4.4).

Atorvastatin

Digoxin: When multiple doses of digoxin and 10 mg atorvastatin were coadministered, steady-state digoxin concentrations increased slightly. Patients taking digoxin should be monitored appropriately.

Oral contraceptives: Coadministration of atorvastatin with an oral contraceptive produced increases in plasma concentrations of norethisterone and ethinyl estradiol.

Warfarin: In a clinical study in patients receiving chronic warfarin therapy, coadministration of atorvastatin 80 mg daily with warfarin caused a small decrease of about 1.7 seconds in prothrombin time during the first 4 days of dosing, which returned to normal within 15 days of atorvastatin treatment.

Although only very rare cases of clinically significant anticoagulant interactions have been reported, prothrombin time should be determined before starting atorvastatin in patients taking coumarin anticoagulants and frequently enough during early therapy to ensure that no significant alteration of prothrombin time occurs. Once a stable prothrombin time has been documented, prothrombin times can be monitored at the intervals usually recommended for patients on coumarin anticoagulants. If the dose of atorvastatin is changed or discontinued, the same procedure should be repeated. Atorvastatin therapy has not been associated with bleeding or with changes in prothrombin time in patients not taking anticoagulants.

Table 1: Effect of co-administered medicinal products on the pharmacokinetics of atorvastatin

Co-administered medicinal product and dosing regimen	Atorvastatin		
	Dose (mg)	Ratio of AUC ^{&}	Clinical Recommendation [#]
Tipranavir 500 mg BID/ Ritonavir 200 mg BID, 8 days (days 14 to 21)	40 mg on day 1, 10 mg on day 20	9.4	In cases where coadministration with atorvastatin is necessary, do not exceed 10 mg atorvastatin daily. Clinical monitoring of these patients is recommended.
Ciclosporin 5.2 mg/kg/day, stable dose	10 mg OD for 28 days	8.7	
Lopinavir 400 mg BID/ Ritonavir 100 mg BID, 14 days	20 mg OD for 4 days	5.9	In cases where coadministration with atorvastatin is necessary, lower maintenance doses of atorvastatin are recommended. At atorvastatin doses exceeding 20 mg, clinical monitoring of these patients is recommended.
Clarithromycin 500 mg BID, 9 days	80 mg OD for 8 days	4.4	
Saquinavir 400 mg BID/ Ritonavir (300 mg BID from days 5-7, increased to 400 mg BID on day 8), days 5-18, 30 min after atorvastatin dosing	40 mg OD for 4 days	3.9	In cases where coadministration with atorvastatin is necessary, lower

Co-administered medicinal product and dosing regimen	Atorvastatin		
	Dose (mg)	Ratio of AUC ^{&}	Clinical Recommendation [#]
Darunavir 300 mg BID/ Ritonavir 100 mg BID, 9 days	10 mg OD for 4 days	3.3	maintenance doses of atorvastatin are recommended. At atorvastatin doses exceeding 40 mg, clinical monitoring of these patients is recommended.
Itraconazole 200 mg OD, 4 days	40 mg SD	3.3	
Fosamprenavir 700 mg BID/ Ritonavir 100 mg BID, 14 days	10 mg OD for 4 days	2.5	
Fosamprenavir 1400 mg BID, 14 days	10 mg OD for 4 days	2.3	
Nelfinavir 1250 mg BID, 14 days	10 mg OD for 28 days	1.74 [^]	No specific recommendation.
Elbasvir 50 mg OD/ Grazoprevir 200 mg OD, 13 days	10 mg SD	1.94	The dose of atorvastatin should not exceed a daily dose of 20 mg during co-administration with products containing elbasvir or grazoprevir.
Glecaprevir 400 mg OD/ Pibrentasvir 120 mg OD, 7 days	10 mg OD for 7 days	8.3	Co-administration with products containing glecaprevir or pibrentasvir is contraindicated (see section 4.3)
Grapefruit Juice, 240 mL OD *	40 mg, SD	1.37	Concomitant intake of large quantities of grapefruit juice and atorvastatin is not recommended.
Diltiazem 240 mg OD, 28 days	40 mg, SD	1.51	After initiation or following dose adjustments of diltiazem, appropriate clinical monitoring of these patients is recommended.
Erythromycin 500 mg QID, 7 days	10 mg, SD	1.33 [^]	Lower maximum dose and clinical monitoring of these patients is recommended.
Amlodipine 10 mg, single dose	80 mg, SD	1.18	No specific recommendation.
Cimetidine 300 mg QID, 2 weeks	10 mg OD for 2 weeks	less than 1.00 [^]	No specific recommendation.
Antacid suspension of magnesium and aluminium hydroxides, 30 mL QID, 2 weeks	10 mg OD for 4 weeks	0.65 [^]	No specific recommendation.
Efavirenz 600 mg OD, 14 days	10 mg for 3 days	0.59	No specific recommendation.
Rifampin 600 mg OD, 7 days (coadministered)	40 mg SD	1.30	If co-administration cannot be avoided, simultaneous

Co-administered medicinal product and dosing regimen	Atorvastatin		
	Dose (mg)	Ratio of AUC ^{&}	Clinical Recommendation [#]
Rifampin 600 mg OD, 5 days (doses separated)	40 mg SD	0.20	coadministration of atorvastatin with rifampin is recommended, with clinical monitoring.
Gemfibrozil 600 mg BID, 7 days	40 mg SD	1.35	Not recommended.
Fenofibrate 160 mg OD, 7 days	40 mg SD	1.03	Not recommended.
Boceprevir 800 mg TID, 7 days	40 mg SD	2.3	Lower starting dose and clinical monitoring of these patients is recommended. The dose of atorvastatin should not exceed a daily dose of 20 mg during co-administration with boceprevir.

[&] Represents ratio of treatments (co-administered drug plus atorvastatin versus atorvastatin alone).

[#] See sections 4.4 and 4.5 for clinical significance.

* Contains one or more components that inhibit CYP3A4 and can increase plasma concentrations of medicinal products metabolised by CYP3A4. Intake of one 240 ml glass of grapefruit juice also resulted in a decreased AUC of 20.4% for the active orthohydroxy metabolite. Large quantities of grapefruit juice (over 1.2 L daily for 5 days) increased AUC of atorvastatin 2.5-fold and AUC of active (atorvastatin and metabolites).

[^] Total atorvastatin equivalent activity

OD = once daily; SD = single dose; BID = twice daily; TID = three times daily; QID = four times daily.

Table 2: Effect of atorvastatin on the pharmacokinetics of co-administered medicinal products

Atorvastatin and dosing regimen	Co-administered medicinal product		
	Medicinal product/Dose (mg)	Ratio of AUC ^{&}	Clinical Recommendation
80 mg OD for 10 days	Digoxin 0.25 mg OD, 20 days	1.15	Patients taking digoxin should be monitored appropriately.
40 mg OD for 22 days	Oral contraceptive OD, 2 months - norethindrone 1 mg - ethinyl estradiol 35 µg	1.28 1.19	No specific recommendation.
80 mg OD for 15 days	* Phenazone, 600 mg SD	1.03	No specific recommendation.
10 mg, OD for 4 days	Fosamprenavir 1400 mg BID, 14 days	0.73	No specific recommendation.

& Represents ratio of treatments (co-administered drug plus atorvastatin versus atorvastatin alone).

* Co-administration of multiple doses of atorvastatin and phenazone showed little or no detectable effect in the clearance of phenazone.

OD = once daily; SD = single dose; BID = twice daily.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women of childbearing potential should use appropriate contraceptive measures during treatment (see section 4.3).

Pregnancy

Ezetimibe/Atorvastatine GL is contraindicated during pregnancy (see section 4.3). No clinical data are available on the use of Ezetimibe/Atorvastatine GL during pregnancy.

Ezetimibe/Atorvastatine GL should not be used in women who are pregnant, trying to become pregnant or suspect they are pregnant. Treatment with Ezetimibe/Atorvastatine GL should be suspended for the duration of pregnancy or until it has been determined that the woman is not pregnant (see section 4.3).

Atorvastatin

Safety in pregnant women has not been established. No controlled clinical trials with atorvastatin have been conducted in pregnant women. Rare reports of congenital anomalies following intrauterine exposure to HMG-CoA reductase inhibitors have been received. Animal studies have shown toxicity to reproduction (see section 5.3).

Maternal treatment with atorvastatin may reduce the fetal levels of mevalonate which is a precursor of cholesterol biosynthesis. Atherosclerosis is a chronic process, and ordinarily discontinuation of lipid-lowering medicinal products during pregnancy should have little impact on the long-term risk associated with primary hypercholesterolaemia.

Ezetimibe

No clinical data are available on the use of ezetimibe during pregnancy. Animal studies on the use of ezetimibe in monotherapy have shown no evidence of direct or indirect harmful effects on pregnancy, embryofoetal development, birth or postnatal development (see section 5.3).

The coadministration of ezetimibe and atorvastatin in pregnant rats indicated that there was a test article-related increase in the skeletal variation “reduced ossification of the sternbrae” in the high dose ezetimibe/atorvastatin group. This may be related to the observed decrease in foetal body weights. In pregnant rabbits a low incidence of skeletal deformities (fused sternbrae, fused caudal vertebrae and asymmetrical sternbrae variations) were observed.

Breastfeeding

Ezetimibe/Atorvastatine GL is contraindicated during breast-feeding.

Because of the potential for serious adverse reactions, women taking Ezetimibe/Atorvastatine GL should not breast-feed their infants (see section 4.3).

Studies on rats have shown that ezetimibe is secreted into breast milk. In rats, plasma concentrations of atorvastatin and its active metabolites are similar to those in milk (see section 5.3). It is not known if the active components of Ezetimibe/Atorvastatine GL are secreted into human breast milk (see section 4.3).

Fertility

No fertility studies were conducted with Ezetimibe/Atorvastatine GL.

Atorvastatin

In animal studies atorvastatin had no effect on male or female fertility.

Ezetimibe

Ezetimibe had no effect on the fertility of male or female rats.

4.7 Effects on ability to drive and use machines

Ezetimibe/Atorvastatine GL has negligible influence on the ability to drive and use machines. However, when driving vehicles or operating machines, it should be taken into account that dizziness has been reported.

4.8 Undesirable effects

Summary of the safety profile

The combination product (or co-administration of ezetimibe and atorvastatin equivalent to the combination product) has been evaluated for safety in more than 2,400 patients in 7 clinical trials.

Tabulated list of adverse reactions

Adverse reactions observed in clinical studies of the combination product (or co-administration of ezetimibe and atorvastatin equivalent to the combination product) or ezetimibe or atorvastatin or reported from postmarketing use with the combination product or ezetimibe or atorvastatin are listed in Table 3. These reactions are presented by system organ class and by frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100, < 1/10$); uncommon ($\geq 1/1000, < 1/100$); rare ($\geq 1/10,000, < 1/1000$); very rare ($< 1/10,000$) and not known (cannot be estimated from the available data).

Table 3: Adverse reactions

System organ class Frequency	Adverse reaction
Infections and infestations	
Uncommon	influenza
Not known	nasopharyngitis
Blood and lymphatic system disorders	
Not known	thrombocytopenia
Immune system disorders	
Not known	hypersensitivity, including anaphylaxis, angioedema, rash, and urticaria
Metabolism and nutrition disorders	
Not known	decreased appetite, anorexia, hyperglycaemia, hypoglycaemia
Psychiatric disorders	
Uncommon	depression, insomnia, sleep disorder
Not known	nightmares
Nervous system disorders	
Uncommon	dizziness, dysgeusia, headache, paraesthesia
Not known	hypoesthesia, amnesia, peripheral neuropathy, myasthenia gravis
Eye disorders	
Not known	vision blurred, visual disturbance, ocular myasthenia
Ear and labyrinth disorders	
Not known	tinnitus, hearing loss
Cardiac disorders	

System organ class Frequency	Adverse reaction
Uncommon	sinus bradycardia
Vascular disorders	
Uncommon	hot flush
Not known	hypertension
Respiratory, thoracic and mediastinal disorders	
Uncommon	dyspnoea
Not known	cough, pharyngolaryngeal pain, epistaxis
Gastrointestinal disorders	
Common	diarrhoea
Uncommon	abdominal discomfort, abdominal distension, abdominal pain, abdominal pain lower, abdominal pain upper, constipation, dyspepsia, flatulence, frequent bowel movements, gastritis, nausea, stomach discomfort
Not known	pancreatitis, gastro-oesophageal reflux disease, eructation, vomiting, dry mouth
Hepatobiliary disorders	
Not known	hepatitis, cholelithiasis, cholecystitis, cholestasis, fatal and non-fatal hepatic failure
Skin and subcutaneous tissue disorders	
Uncommon	acne, urticaria
Not known	alopecia, skin rash, pruritus, erythema multiforme, angioedema, dermatitis bullous including erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis
Musculoskeletal and connective tissue disorders	
Common	myalgia
Uncommon	arthralgia, back pain, muscle fatigue, muscle spasms, muscular weakness, pain in extremity
Not known	myopathy/rhabdomyolysis, muscle rupture, tendinopathy sometimes complicated by rupture, neck pain, joint swelling, myositis, lupus-like syndrome, immune-mediated necrotizing myopathy (see section 4.4)
Reproductive system and breast disorders	
Not known	gynaecomastia
General disorders and administration site conditions	
Uncommon	asthenia, fatigue, malaise, oedema
Not known	chest pain, pain, peripheral oedema, pyrexia
Investigations	
Uncommon	ALT and/or AST increased, alkaline phosphatase increased, blood creatine phosphokinase (CPK) increased, gamma-glutamyltransferase increased, hepatic enzyme increased, liver function test abnormal, weight increased
Not known	white blood cells urine positive

Laboratory Values

In controlled clinical trials, the incidence of clinically important elevations in serum transaminases (ALT and/or AST ≥ 3 X ULN, consecutive) was 0.6% for patients treated with the combination

product. These elevations were generally asymptomatic, not associated with cholestasis, and returned to baseline spontaneously or after discontinuation of therapy (see section 4.4).

The following adverse events have been reported with some statins:

- sexual dysfunction
- depression
- exceptional cases of interstitial lung disease, especially with long term therapy (see section 4.4)
- diabetes mellitus: frequency will depend on the presence or absence of risk factors (fasting blood glucose ≥ 5.6 mmol/L, BMI > 30 kg/m², raised triglycerides, history of hypertension)

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

4.9 Overdose

In the event of an overdose, symptomatic and supportive measures should be employed. Liver function tests should be performed and serum CPK levels should be monitored.

Ezetimibe

In clinical studies, administration of ezetimibe, 50 mg/day to 15 healthy subjects for up to 14 days, or 40 mg/day to 18 patients with primary hypercholesterolaemia for up to 56 days, was generally well tolerated. A few cases of overdose have been reported; most have not been associated with adverse experiences. Reported adverse experiences have not been serious. In animals, no toxicity was observed after single oral doses of 5000 mg/kg of ezetimibe in rats and mice and 3000 mg/kg in dogs.

Atorvastatin

Due to extensive atorvastatin binding to plasma proteins, haemodialysis is not expected to significantly enhance atorvastatin clearance.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Lipid modifying agents, HMG-CoA reductase inhibitors in combination with other lipid modifying agents, ATC code: C10BA05

Ezetimibe/Atorvastatine GL (ezetimibe/atorvastatin) is a lipid-lowering product that selectively inhibits the intestinal absorption of cholesterol and related plant sterols and inhibits the endogenous synthesis of cholesterol.

Mechanism of action

Plasma cholesterol is derived from intestinal absorption and endogenous synthesis. Ezetimibe/Atorvastatine GL contains ezetimibe and atorvastatin, two lipid lowering compounds with complementary mechanisms of action. Ezetimibe/Atorvastatine GL reduces elevated total cholesterol (total-C), LDL-C, apolipoprotein B (Apo B), triglycerides (TG), and non-high-density lipoprotein cholesterol (non-HDL-C), and increases high-density lipoprotein cholesterol (HDL-C) through dual inhibition of cholesterol absorption and synthesis.

Ezetimibe

Ezetimibe inhibits the intestinal absorption of cholesterol. Ezetimibe is orally active and has a mechanism of action that differs from other classes of cholesterol-reducing compounds (e.g. statins, bile acid sequestrants [resins], fibric acid derivatives, and plant stanols). The molecular target of ezetimibe is the sterol transporter, Niemann-Pick C1 Like 1 (NPC1L1), which is responsible for the intestinal uptake of cholesterol and phytosterols.

Ezetimibe localises at the brush border of the small intestine and inhibits the intestinal absorption of cholesterol, leading to a decrease in the delivery of intestinal cholesterol to the liver; statins reduce cholesterol synthesis in the liver and together these distinct mechanisms provide complementary cholesterol reduction. In a 2-week clinical study in 18 hypercholesterolaemic patients, ezetimibe inhibited intestinal cholesterol absorption by 54%, compared with placebo.

A series of preclinical studies were performed to determine the selectivity of ezetimibe for inhibiting cholesterol absorption. Ezetimibe inhibited the absorption of [¹⁴C]-cholesterol with no effect on the absorption of triglycerides, fatty acids, bile acids, progesterone, ethinyl estradiol, or fat soluble vitamins A and D.

Atorvastatin

Atorvastatin is a selective, competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme responsible for the conversion of 3-hydroxy-3-methyl-glutaryl-coenzyme A to mevalonate, a precursor of sterols, including cholesterol. Triglycerides and cholesterol in the liver are incorporated into very low-density lipoproteins (VLDL) and released into the plasma for delivery to peripheral tissues. Low-density lipoproteins (LDL) are formed from VLDL and are catabolized primarily through the receptor with high affinity to LDL (LDL receptor).

Atorvastatin lowers plasma cholesterol and lipoprotein serum concentrations by inhibiting HMG-CoA reductase and subsequently cholesterol biosynthesis in the liver and increases the number of hepatic LDL receptors on the cell surface for enhanced uptake and catabolism of LDL.

Atorvastatin reduces LDL production and the number of LDL particles. Atorvastatin produces a profound and sustained increase in LDL receptor activity coupled with a beneficial change in the quality of circulating LDL particles. Atorvastatin is effective in reducing LDL-C in patients with homozygous familial hypercholesterolaemia, a population that has not usually responded to lipid-lowering medicinal products.

Atorvastatin has been shown to reduce concentrations of total-C (30%-46%), LDL-C (41%-61%), apolipoprotein B (34%-50%), and triglycerides (14%-33%) while producing variable increases in HDL-C and apolipoprotein A1 in a dose response study. These results are consistent in patients with heterozygous familial hypercholesterolaemia, nonfamilial forms of hypercholesterolaemia, and mixed hyperlipidaemia, including patients with noninsulin-dependent diabetes mellitus.

Clinical efficacy and safety

In controlled clinical studies, the combination product significantly reduced total-C, LDL-C, Apo B, and TG, and increased HDL-C in patients with hypercholesterolaemia.

In a placebo-controlled study, 628 patients with hyperlipidaemia were randomised to receive placebo, ezetimibe (10 mg), atorvastatin (10 mg, 20 mg, 40 mg, or 80 mg), or coadministered ezetimibe and atorvastatin equivalent to Atorvastatin/Ezetimibe (10/10, 10/20, 10/40 and 10/80) for up to 12 weeks. Patients receiving all doses of Atorvastatin/Ezetimibe were compared to those receiving all doses of atorvastatin. Atorvastatin/Ezetimibe lowered total-C, LDL-C, Apo B, TG, and non-HDL-C, and increased HDL-C significantly more than atorvastatin alone.

In a multicenter, randomised, double-blind study, 621 adults patients with HeFH, coronary heart disease, or multiple (≥ 2) cardiovascular risk factors, and a LDL-C level ≥ 130 mg/dL received after a

6- to 10-week dietary stabilization and atorvastatin (10 mg/day) open-label run-in period atorvastatin 10 mg + ezetimibe 10 mg or atorvastatin 20 mg. Atorvastatin dose in both groups was doubled after 4 weeks, 9 weeks, or both when the LDL-C level was not at its goal ≤ 100 mg/dL) to a maximum of 40 mg in the combination group and 80 mg in the monotherapy group. The proportion of subjects reaching their target LDL-C level goal of ≤ 100 mg/dL was significantly higher in the combination group than in the atorvastatin monotherapy group (22% vs 7%; $P < 0.01$). At 4 weeks, levels of LDL-C, triglycerides, and non-high-density lipoprotein cholesterol were reduced significantly more by combination therapy than by doubling the dose of atorvastatin (LDL-C -22.8% versus -8.6%; $P < 0.01$).

In another randomised, double-blind, placebo-controlled study, 450 hypercholesterolaemic patients with coronary heart disease who had not achieved their LDL-C goal ≤ 2.60 mmol/L while on a stable dose of atorvastatin 10 or 20 mg/day for ≥ 6 weeks received either atorvastatin + ezetimibe or atorvastatin + placebo. Significantly more patients achieved an LDL-C goal ≤ 2.6 mmol/L with ezetimibe than placebo (81.3 vs. 21.8%; $P \leq 0.001$). Compared to placebo, co-administration of ezetimibe with ongoing atorvastatin led to significantly ($P \leq 0.001$) greater reductions in LDL-C, TC, TG, non-HDL-C, and apolipoprotein B; HDL-C was significantly ($P \leq 0.05$) increased.

5.2 Pharmacokinetic properties

The combination product has been shown to be bioequivalent to coadministration of corresponding doses of ezetimibe and atorvastatin tablets.

Absorption

The effects of a high-fat meal on the pharmacokinetics of ezetimibe and atorvastatin when administered as combination product are comparable to those reported for the individual tablets.

Ezetimibe

After oral administration, ezetimibe is rapidly absorbed and extensively conjugated to a pharmacologically active phenolic glucuronide (ezetimibe-glucuronide). Mean maximum plasma concentrations (C_{max}) occur within 1-2 hours for ezetimibe-glucuronide and 4-12 hours for ezetimibe. The absolute bioavailability of ezetimibe cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection.

Concomitant food administration (high-fat or non-fat meals) had no effect on the oral bioavailability of ezetimibe when administered at 10-mg-tablets.

Atorvastatin

Atorvastatin is rapidly absorbed after oral administration; maximum plasma concentrations (C_{max}) occur within 1-2 hours. Extent of absorption increases in proportion to atorvastatin dose. After oral administration, atorvastatin tablets are 95%-99% bioavailable compared to the oral solution. The absolute bioavailability of atorvastatin is approximately 12% and the systemic availability of HMG-CoA reductase inhibitory activity is approximately 30%. The low systemic availability is attributed to presystemic clearance in gastrointestinal mucosa and/or hepatic first-pass metabolism.

Distribution

Ezetimibe

Ezetimibe and ezetimibe-glucuronide are bound 99.7% and 88-92% to human plasma proteins, respectively.

Atorvastatin

Mean volume of distribution of atorvastatin is approximately 381 L. Atorvastatin is $\geq 98\%$ bound to plasma proteins.

Biotransformation

Ezetimibe

Ezetimibe is metabolised primarily in the small intestine and liver via glucuronide conjugation (a phase II reaction) with subsequent biliary excretion. Minimal oxidative metabolism (a phase I reaction) has been observed in all species evaluated. Ezetimibe and ezetimibe-glucuronide are the major drug-derived compounds detected in plasma, constituting approximately 10-20% and 80-90% of the total drug in plasma, respectively. Both ezetimibe and ezetimibe-glucuronide are slowly eliminated from plasma with evidence of significant enterohepatic recycling. The half-life for ezetimibe and ezetimibe-glucuronide is approximately 22 hours.

Atorvastatin

Atorvastatin is metabolized by cytochrome P450 3A4 to ortho- and parahydroxylated derivatives and various beta-oxidation products. Apart from other pathways these products are further metabolised via glucuronidation. *In vitro*, inhibition of HMG-CoA reductase by ortho- and parahydroxylated metabolites is equivalent to that of atorvastatin. Approximately 70% of circulating inhibitory activity for HMG-CoA reductase is attributed to active metabolites.

Elimination

Ezetimibe

Following oral administration of ¹⁴C-ezetimibe (20 mg) to human subjects, total ezetimibe accounted for approximately 93% of the total radioactivity in plasma. Approximately 78% and 11% of the administered radioactivity were recovered in the faeces and urine, respectively, over a 10-day collection period. After 48 hours, there were no detectable levels of radioactivity in the plasma.

Atorvastatin

Atorvastatin is a substrate of the hepatic transporters, organic anion-transporting polypeptide 1B1 (OATP1B1) and 1B3 (OATP1B3) transporter. Metabolites of atorvastatin are substrates of OATP1B1. Atorvastatin is also identified as a substrate of the efflux transporters multi-drug resistance protein 1 (MDR1) and breast cancer resistance protein (BCRP), which may limit the intestinal absorption and biliary clearance of atorvastatin. Atorvastatin is eliminated primarily in bile following hepatic and/or extrahepatic metabolism. However, the medicinal product does not appear to undergo significant enterohepatic recirculation. Mean plasma elimination half-life of atorvastatin in humans is approximately 14 hours. The half-life of inhibitory activity of HMG-CoA reductase is approximately 20-30 hours due to the contribution of active metabolites.

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with Ezetimibe/Atorvastatin GL in all subsets of the paediatric population.

Ezetimibe

The pharmacokinetics of ezetimibe are similar between children ≥ 6 years and adults. Pharmacokinetic data in the paediatric population < 6 years of age are not available. Clinical experience in paediatric and adolescent patients includes patients with HoFH, HeFH or sitosterolaemia.

Atorvastatin

In an open-label, 8-week study, Tanner Stage 1 (n=15) and Tanner Stage 2 (n=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.

Elderly

Ezetimibe

Plasma concentrations for total ezetimibe are about 2-fold higher in the elderly (≥ 65 years) than in the young (18 to 45 years). LDL-C reduction and safety profile are comparable between elderly and younger subjects treated with ezetimibe.

Atorvastatin

Plasma concentrations of atorvastatin and its active metabolites are higher in healthy elderly subjects than in young adults while the lipid effects were comparable to those seen in younger patient populations.

Hepatic impairment

Ezetimibe

After a single 10-mg dose of ezetimibe, the mean AUC for total ezetimibe was increased approximately 1.7-fold in patients with mild hepatic insufficiency (Child-Pugh score 5 or 6), compared to healthy subjects. In a 14-day, multiple-dose study (10 mg daily) in patients with moderate hepatic insufficiency (Child-Pugh score 7-9), the mean AUC for total ezetimibe was increased approximately 4-fold on Day 1 and Day 14 compared to healthy subjects. No dose adjustment is necessary for patients with mild hepatic insufficiency. Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate or severe (Child-Pugh score > 9) hepatic insufficiency, ezetimibe is not recommended in these patients (see sections 4.2 and 4.4).

Atorvastatin

Plasma concentrations of atorvastatin and its active metabolites are markedly increased (approx. 16-fold in C_{max} and approx. 11-fold in AUC) in patients with chronic alcoholic liver disease (Child-Pugh B).

Renal impairment

Ezetimibe

After a single 10-mg dose of ezetimibe in patients with severe renal disease ($n=8$; mean $CrCl \leq 30$ ml/min/1.73 m²), the mean AUC for total ezetimibe was increased approximately 1.5-fold, compared to healthy subjects ($n=9$).

An additional patient in this study (post-renal transplant and receiving multiple medicinal products, including ciclosporin) had a 12-fold greater exposure to total ezetimibe.

Atorvastatin

Renal disease has no influence on the plasma concentrations or lipid effects of atorvastatin and its active metabolites.

Gender

Ezetimibe

Plasma concentrations for total ezetimibe are slightly higher (approximately 20%) in women than in men. LDL-C reduction and safety profile are comparable between men and women treated with ezetimibe.

Atorvastatin

Concentrations of atorvastatin and its active metabolites in women differ from those in men (women: approx. 20% higher for C_{max} and approx. 10% lower for AUC). These differences were of no clinical significance, resulting in no clinically significant differences in lipid effects among men and women.

SLCO1B1 polymorphism

Atorvastatin

Hepatic uptake of all HMG-CoA reductase inhibitors, including atorvastatin, involves the OATP1B1 transporter. In patients with SLCO1B1 polymorphism there is a risk of increased exposure of atorvastatin, which may lead to an increased risk of rhabdomyolysis (see section 4.4). Polymorphism in the gene encoding OATP1B1 (SLCO1B1 c.521CC) is associated with a 2.4-fold higher atorvastatin exposure (AUC) than in individuals without this genotype variant (c.521TT). A genetically impaired hepatic uptake of atorvastatin is also possible in these patients. Possible consequences for the efficacy are unknown.

5.3 Preclinical safety data

Ezetimibe

Animal studies on the chronic toxicity of ezetimibe identified no target organs for toxic effects. In dogs treated for four weeks with ezetimibe (≥ 0.03 mg/kg/day) the cholesterol concentration in the cystic bile was increased by a factor of 2.5-3.5. However, in a one year study on dogs given doses of up to 300 mg/kg/day no increased incidence of cholelithiasis or other hepatobiliary effects were observed. The significance of these data for humans is not known. A lithogenic risk associated with the therapeutic use of ezetimibe cannot be ruled out. Long-term carcinogenicity tests on ezetimibe were negative. Ezetimibe had no effect on the fertility of male or female rats, nor was it found to be teratogenic in rats or rabbits, nor did it affect prenatal or postnatal development. Ezetimibe crossed the placental barrier in pregnant rats and rabbits given multiple doses of 1000 mg/kg/day.

Atorvastatin

Atorvastatin was negative for mutagenic and clastogenic potential in a battery of 4 *in vitro* tests and 1 *in vivo* assay. Atorvastatin was not found to be carcinogenic in rats, but high doses in mice (resulting in 6- to 11-fold the AUC_{0-24h} reached in humans at the highest recommended dose) showed hepatocellular adenomas in males and hepatocellular carcinomas in females. There is evidence from animal experimental studies that HMG-CoA reductase inhibitors may affect the development of embryos or fetuses. In rats, rabbits, and dogs atorvastatin had no effect on fertility and was not teratogenic, however, at maternally toxic doses foetal toxicity was observed in rats and rabbits. The development of the rat offspring was delayed and post-natal survival reduced during exposure of the dams to high doses of atorvastatin. In rats, there is evidence of placental transfer. In rats, plasma concentrations of atorvastatin are similar to those in milk. It is not known whether atorvastatin or its metabolites are excreted in human milk.

Ezetimibe and statin coadministered

In three-month co-administration studies in rats and dogs with ezetimibe and atorvastatin, the toxic effects observed were essentially those typically associated with statins. The statin-like histopathologic findings were limited to the liver. Some of the toxic effects were more pronounced than observed during treatment with statins alone. This is attributed to pharmacokinetic and pharmacodynamic interactions in co-administration therapy.

The co-administration of ezetimibe and atorvastatin in pregnant rats indicated that there was a test article-related increase in the skeletal variation “reduced ossification of the sternbrae” in the high dose (1,000/108.6 mg/kg) ezetimibe/atorvastatin group. This may be related to the observed decrease in foetal body weights. In pregnant rabbits a low incidence of skeletal deformities (fused sternbrae, fused caudal vertebrae and asymmetrical sternbrae variation) were observed.

In a series of *in vivo* and *in vitro* assays, ezetimibe, given alone or co-administered with atorvastatin, exhibited no genotoxic potential.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Lactose monohydrate
Calcium carbonate
Cellulose, microcrystalline
Sodium Lauryl Sulfate [E487]
Croscarmellose sodium
Povidone K30
Hydroxypropylcellulose
Magnesium Stearate
Polysorbate 80

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

2 years.

6.4 Special precautions for storage

Store in the original package in order to protect from light.

6.5 Nature and contents of container

Ezetimibe/Atorvastatine GL 10 mg/10, 20, 40, 80 mg tabletten will be available in packs of 10, 30, 90 and 100 tablets in an OPA/Aluminium/PVC blister in a carton.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. HOUDER VAN DE VERGUNNING VOOR HET IN DE HANDEL BRENGEN

G.L. Pharma GmbH
Schlossplatz 1
8502 Lannach
Oostenrijk

8. NUMMER(S) VAN DE VERGUNNING VOOR HET IN DE HANDEL BRENGEN

Ezetimibe/Atorvastatine GL 10 mg/ 10 mg, tabletten	RVG 125040
Ezetimibe/Atorvastatine GL 10 mg/ 20 mg, tabletten	RVG 125041

Ezetimibe/Atorvastatine GL 10 mg/ 40 mg, tabletten RVG 125042
Ezetimibe/Atorvastatine GL 10 mg/ 80 mg, tabletten RVG 125043

9. DATUM VAN EERSTE VERLENING VAN DE VERGUNNING/VERLENGING VAN DE VERGUNNING

Datum van eerste verlening van de vergunning: 21 juli 2020

10. DATUM VAN HERZIENING VAN DE TEKST

Laatst gedeeltelijke wijziging betreft de rubrieken 4.4, en 4.8: 14 maart 2023.