

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

Glytrin, 400 microgram per dosis, sublinguale spray

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Glyceryl Trinitrate: 400 micrograms per metered dose

Excipient(s) with known effect

Ethanol (alcohol) – 7.5 mg per spray.

For the full list of excipients, see Section 6.1

### 3. PHARMACEUTICAL FORM

Sublingual spray, solution.

(Sublingual spray)

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Treatment of acute angina pectoris.

Prevention of inducible angina (e.g. physical effort, emotional stress, exposure to cold).

#### 4.2 Posology and method of administration

Posology

Oromucosal Dosage

*Adults including the elderly*

At the onset of an attack, one or two metered doses (400 to 800 micrograms glyceryl trinitrate) to be sprayed under the tongue for the relief of anginal pain while breath is held. No more than three doses are recommended at any one time.

For the prevention of inducible angina (e.g. physical effort, emotional stress, exposure to cold), one or two 400 microgram metered doses sprayed under the tongue within 2 – 3 minutes of the event starting.

*Paediatric population*

Glytrin should not be used in children and adolescents under 18 years.

Method of administration

Before using Glytrin for the first time, the patient should check that the spray is working by pressing the pump button a few times until it produces a fine mist of liquid. The patient should practice aiming the spray onto a tissue or similar item so that they will be able to aim it correctly under the tongue when they need to use it. If the patient does not need to use Glytrin very often, the spray should be checked regularly to see that it still works properly.

During application the patient should rest, ideally in the sitting position. The canister should be held vertically with the valve head uppermost and the spray orifice as close to the mouth as possible. The dose should be sprayed under the tongue and the mouth should be closed immediately after each dose. The spray should not

be inhaled. Patients should be instructed to familiarise themselves with the position of the spray orifice, which can be identified by the finger rest on top of the valve, in order to facilitate orientation, for administration at night.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- Severe hypotension (systolic blood pressure lower than 90 mm Hg)
- Hypotensive shock
- Severe anaemia
- Constrictive pericarditis
- Extreme bradycardia
- Glucose-6-phosphate-dehydrogenase-deficiency
- Cerebral haemorrhage and brain trauma
- Aortic and/or mitral stenosis
- Angina caused by hypertrophic obstructive cardiomyopathy
- Circulatory collapse
- Cardiogenic shock
- Toxic pulmonary oedema
- Concomitant use with phosphodiesterase inhibitors, such as sildenafil, tadalafil, or vardenafil
- Concomitant use with the soluble guanylate cyclase stimulator riociguat (see section 4.5).

### **4.4 Special warnings and precautions for use**

Tolerance to this drug and cross-tolerance to other nitrates may occur.

Glytrin should be administered with particular caution in:

- Pericardial tamponade
- Low filling pressures (e.g. acute myocardial infarction, left ventricular failure)
- Tendency to dysregulation of orthostatic blood pressure
- Diseases accompanied by an increase in intracranial pressure (so far further pressure has been observed solely in high doses of glyceryl trinitrate).

Alcohol should be avoided because of the hypotensive effect and medical controls of the intraocular pressure of glaucoma patients are advisable.

Particular caution should also be exercised when using Glytrin in patients with volume depletion from diuretic therapy, severe hepatic or renal impairment and hypothyroidism.

This medicine contains 7.5 mg of alcohol (ethanol) in each spray. The amount in one spray of this medicine is equivalent to less than 1 ml beer or 1 ml wine. The small amount of alcohol in this medicine will not have any noticeable effects.

### **4.5 Interaction with other medicinal products and other forms of interaction**

- Alcohol may potentiate the hypotensive effect.
- Vasodilators, antihypertensives,  $\beta$ -blockers, calcium antagonists, neuroleptics, tricyclic antidepressants and diuretics can increase nitrate-induced hypotension.
- The hypotensive effects of nitrates are potentiated by the concurrent administration of phosphodiesterase inhibitors, such as sildenafil, tadalafil, or vardenafil.
- The use of a soluble guanylate cyclase stimulator such as riociguat is contraindicated (see section 4.3) since concomitant use can cause hypotension.
- The bioavailability of dihydroergotamine may be increased by concomitant use of Glytrin, which can result in vasoconstriction since dihydroergotamine can antagonise the effects of glyceryl trinitrate.

- The concomitant administration of Glytrin and heparin can reduce the antithrombotic effect of heparin. Regular monitoring of coagulation parameters and adjustment of the heparin dose may be necessary.
- In patients pre-treated with organic nitrates a higher dose of glyceryl trinitrate may be necessary to achieve the desired haemodynamic effect.

#### 4.6 Fertility, pregnancy and lactation

##### Pregnancy

The safety of glyceryl trinitrate in human pregnancy, especially during the first trimester has not been established.

##### Breastfeeding

It is not known whether glyceryl trinitrate is excreted into human breast milk. Glytrin should be used only after weighing the benefit for the mother against possible risks for the child. Nursing should be discontinued during treatment with this product.

##### Fertility

Preclinical data reveal no special hazard for humans based on conventional studies of toxicity to reproduction (see section 5.3).

#### 4.7 Effects on ability to drive and use machines

The ability to react may be diminished because of the side effects or interactions due to the nitrates. This effect is potentiated by alcohol consumption. Therefore, driving and/or using machines should be avoided during treatment with Glytrin.

#### 4.8 Undesirable effects

The following adverse reactions have been reported:

System Organ Class	Very common (≥ 1/10)	Common (≥ 1/100 to < 1/10)	Uncommon (≥ 1/1,000 to < 1/100)	Rare (≥ 1/10,000 to < 1/1,000)	Very Rare (< 1/10,000)
<b>Nervous System Disorders</b>	Headache	Vertigo Dizziness		Syncope	
<b>Skin and Subcutaneous Tissue Disorders</b>				Allergic dermatitis *	Exfoliative dermatitis
<b>Vascular Disorders</b>		Facial flushing		Orthostatic hypotension Circulatory collapse	
<b>General Disorders and Administration Site Conditions</b>		Asthenia	Application site discomfort including Burning sensation and Stinging		
<b>Gastrointestinal Disorders</b>		Nausea	Tongue blistering		

System Organ Class	Very common (≥ 1/10)	Common (≥ 1/100 to < 1/10)	Uncommon (≥ 1/1,000 to < 1/100)	Rare (≥ 1/10,000 to < 1/1,000)	Very Rare (< 1/10,000)
<b>Cardiac Disorders</b>				Tachycardia Bradycardia Angina pectoris aggravated	
<b>Investigations</b>				Blood pressure decreased	

\* symptoms which are known in conjunction with hypersensitivity reactions

Use of Glytrin may give rise to transient hypoxaemia and, in patients with coronary heart disease, ischaemia as a result of a relative redistribution of the bloodstream, which is to hypoventilated alveolar areas.

Tolerance development and the occurrence of crossed tolerance of other nitro compounds have been found in chronic, continuous treatment using high doses. To avoid a decrease in efficacy or a loss of efficacy, high continuous doses should be avoided.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Nederlands Bijwerkingen Centrum Lareb Website: [www.lareb.nl](http://www.lareb.nl).

## 4.9 Overdose

#### Signs and symptoms

Flushing, severe headache, vertigo, tachycardia, a feeling of suffocation, hypotension, fainting and rarely cyanosis and methaemoglobinaemia may occur. In a few patients, there may be a reaction comparable to shock with nausea, vomiting, weakness, sweating and syncope.

#### Treatment

Recovery often occurs without special treatment. Hypotension may be corrected by elevation of the legs to promote venous return. Methaemoglobinaemia should be treated by intravenous methylthioninium chloride and / or toluidine blue. Symptomatic treatment should be given for respiratory and circulatory defects in more serious cases.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Vasodilators used in cardiac diseases, organic nitrates.  
ATC-Code: C01DA02

Glyceryl trinitrate acts on vascular smooth muscles to produce arterial and venous vasodilation. The vasodilation results in a reduction of venous return and an improvement in myocardial perfusion with the result of a reduction in the work performed by the heart and hence reduced oxygen demand.

### 5.2 Pharmacokinetic properties

Glyceryl trinitrate is rapidly absorbed through the buccal and sublingual mucosa, and in man peak concentrations in plasma are observed within four minutes of sublingual administration.

The absolute bioavailability after sublingual administration is approximately 39%. After sublingual administration the plasma levels have shown a wide range of intra and inter-individual variability. The compound is extensively metabolised by liver enzymes and has a plasma half-life of 1-3 minutes. The principle mechanism of metabolism involves denitration.

### **5.3 Preclinical safety data**

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, or toxicity to reproduction.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Peppermint oil  
Propellant HFC 134A (1,1,1,2 Tetrafluoroethane)  
Ethanol

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Do not store above 25°C. Do not refrigerate or freeze.

Glytrin is an aerosol spray and contains a pressurised liquid. Do not expose to temperatures higher than 50°C, and do not pierce the aluminium container (canister), even when empty.

It should not be sprayed at a naked flame or any incandescent material.

Patients, especially those who smoke should be warned not to use Glytrin near a naked flame.

### **6.5 Nature and contents of container**

Internally lacquered monobloc aluminium pressurised container (canister), sealed with a metered spray valve. The product is presented in packs with one container.

Each container contains 1760.0 mg of solution (according to 11400.0 mg of solution and propellant) providing 200 single metered doses.

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Evolan Pharma AB  
Svärdvägen 19  
SE-182 33 Danderyd  
Zweden

## **8. MARKETING AUTHORISATION NUMBER**

RVG 21493

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Datum van eerste verlening van de vergunning: 14 december 1998

Datum van laatste verlenging: 3 september 2006

**10. DATE OF REVISION OF THE TEXT**

Laatste gedeeltelijke wijziging betreft rubriek 7: 23 december 2024.